

# Sum theorems for maximally monotone operators of type (FPV)

Jonathan M. Borwein\* and Liangjin Yao†

May 29, 2013

## Abstract

The most important open problem in Monotone Operator Theory concerns the maximal monotonicity of the sum of two maximally monotone operators provided that the classical Rockafellar's constraint qualification holds.

In this paper, we establish the maximal monotonicity of  $A + B$  provided that  $A$  and  $B$  are maximally monotone operators such that  $\text{star}(\text{dom } A) \cap \text{int } \text{dom } B \neq \emptyset$ , and  $A$  is of type (FPV). We show that when also  $\text{dom } A$  is convex, the sum operator:  $A + B$  is also of type (FPV). Our result generalizes and unifies several recent sum theorems.

## 2010 Mathematics Subject Classification:

Primary 47H05; Secondary 49N15, 52A41, 90C25

**Keywords:** Constraint qualification, convex function, convex set, Fitzpatrick function, linear relation, maximally monotone operator, monotone operator, operator of type (FPV), sum problem.

## 1 Introduction

Throughout this paper, we assume that  $X$  is a real Banach space with norm  $\|\cdot\|$ , that  $X^*$  is the continuous dual of  $X$ , and that  $X$  and  $X^*$  are paired by  $\langle \cdot, \cdot \rangle$ . Let  $A: X \rightrightarrows X^*$  be a *set-valued operator* (also known as a relation, point-to-set mapping or multifunction) from  $X$  to  $X^*$ , i.e., for every  $x \in X$ ,  $Ax \subseteq X^*$ , and let  $\text{gra } A := \{(x, x^*) \in X \times X^* \mid x^* \in Ax\}$  be the *graph* of  $A$ . Recall that  $A$  is *monotone* if

$$\langle x - y, x^* - y^* \rangle \geq 0, \quad \forall (x, x^*) \in \text{gra } A \quad \forall (y, y^*) \in \text{gra } A,$$

---

\*CARMA, University of Newcastle, Newcastle, New South Wales 2308, Australia. E-mail: jonathan.borwein@newcastle.edu.au. Laureate Professor at the University of Newcastle and Distinguished Professor at King Abdul-Aziz University, Jeddah.

†CARMA, University of Newcastle, Newcastle, New South Wales 2308, Australia. E-mail: liangjin.yao@newcastle.edu.au.

and *maximally monotone* if  $A$  is monotone and  $A$  has no proper monotone extension (in the sense of graph inclusion). Let  $A : X \rightrightarrows X^*$  be monotone and  $(x, x^*) \in X \times X^*$ . We say  $(x, x^*)$  is *monotonically related to*  $\text{gra } A$  if

$$\langle x - y, x^* - y^* \rangle \geq 0, \quad \forall (y, y^*) \in \text{gra } A.$$

Let  $A : X \rightrightarrows X^*$  be maximally monotone. We say  $A$  is *of type (FPV)* if for every open convex set  $U \subseteq X$  such that  $U \cap \text{dom } A \neq \emptyset$ , the implication

$$x \in U \text{ and } (x, x^*) \text{ is monotonically related to } \text{gra } A \cap (U \times X^*) \Rightarrow (x, x^*) \in \text{gra } A$$

holds. We emphasize that it remains possible that all maximally monotone operators are of type (FPV). Also every (FPV) operator has the closure of its domain convex. See [28, 27, 8, 10] for this and more information on operators of type (FPV). We say  $A$  is a *linear relation* if  $\text{gra } A$  is a linear subspace.

Monotone operators have proven important in modern Optimization and Analysis; see, e.g., the books [1, 8, 14, 15, 20, 27, 28, 25, 40, 41, 42] and the references therein. We adopt standard notation used in these books: thus,  $\text{dom } A := \{x \in X \mid Ax \neq \emptyset\}$  is the *domain* of  $A$ . Given a subset  $C$  of  $X$ ,  $\text{int } C$  is the *interior* of  $C$ ,  $\text{bdry } C$  is the *boundary* of  $C$ ,  $\text{aff } C$  is the *affine hull* of  $C$ ,  $\overline{C}$  is the *norm closure* of  $C$ , and  $\text{span } C$  is the *span* (the set of all finite linear combinations) of  $C$ . The *intrinsic core* or *relative algebraic interior* of  $C$ ,  ${}^iC$  [40], is defined by  ${}^iC := \{a \in C \mid \forall x \in \text{aff}(C - C), \exists \delta > 0, \forall \lambda \in [0, \delta] : a + \lambda x \in C\}$ . We then define  ${}^{ic}C$  by

$${}^{ic}C := \begin{cases} {}^iC, & \text{if } \text{aff } C \text{ is closed;} \\ \emptyset, & \text{otherwise,} \end{cases}$$

The *indicator function* of  $C$ , written as  $\iota_C$ , is defined at  $x \in X$  by

$$\iota_C(x) := \begin{cases} 0, & \text{if } x \in C; \\ \infty, & \text{otherwise.} \end{cases}$$

If  $C, D \subseteq X$ , we set  $C - D = \{x - y \mid x \in C, y \in D\}$ . For every  $x \in X$ , the *normal cone* operator of  $C$  at  $x$  is defined by  $N_C(x) := \{x^* \in X^* \mid \sup_{c \in C} \langle c - x, x^* \rangle \leq 0\}$ , if  $x \in C$ ; and  $N_C(x) = \emptyset$ , if  $x \notin C$ . We define the *support points* of  $C$ , written as  $\text{supp } C$ , by  $\text{supp } C := \{c \in C \mid N_C(c) \neq \{0\}\}$ . For  $x, y \in X$ , we set  $[x, y] := \{tx + (1 - t)y \mid 0 \leq t \leq 1\}$ . We define the *centre* or *star* of  $C$  by  $\text{star } C := \{x \in C \mid [x, c] \subseteq C, \forall c \in C\}$  [7]. Then  $C$  is convex if and only if  $\text{star } C = C$ .

Given  $f : X \rightarrow ]-\infty, +\infty]$ , we set  $\text{dom } f := f^{-1}(\mathbb{R})$ . We say  $f$  is *proper* if  $\text{dom } f \neq \emptyset$ . We also set  $P_X : X \times X^* \rightarrow X : (x, x^*) \mapsto x$ . Finally, the *open unit ball* in  $X$  is denoted by  $U_X := \{x \in X \mid \|x\| < 1\}$ , the *closed unit ball* in  $X$  is denoted by  $B_X := \{x \in X \mid \|x\| \leq 1\}$ , and  $\mathbb{N} := \{1, 2, 3, \dots\}$ . We denote by  $\rightarrow$  and  $\rightarrow_{w^*}$  the norm convergence and weak\* convergence of nets, respectively.

Let  $A$  and  $B$  be maximally monotone operators from  $X$  to  $X^*$ . Clearly, the *sum operator*  $A + B : X \rightrightarrows X^* : x \mapsto Ax + Bx := \{a^* + b^* \mid a^* \in Ax \text{ and } b^* \in Bx\}$  is monotone. Rockafellar established the following very important result in 1970.

**Theorem 1.1 (Rockafellar’s sum theorem)** (See [24, Theorem 1] or [8].) *Suppose that  $X$  is reflexive. Let  $A, B : X \rightrightarrows X^*$  be maximally monotone. Assume that  $A$  and  $B$  satisfy the classical constraint qualification*

$$\text{dom } A \cap \text{int dom } B \neq \emptyset.$$

*Then  $A + B$  is maximally monotone.*

Arguably, the most significant open problem in the theory concerns the maximal monotonicity of the sum of two maximally monotone operators in general Banach spaces; this is called the “sum problem”. Some recent developments on the sum problem can be found in Simons’ monograph [28] and [4, 5, 6, 8, 12, 11, 36, 19, 31, 37, 38, 39]. It is known, among other things, that the sum theorem holds under Rockafellar’s constraint qualification when both operators are of dense type or when each operator has nonempty domain interior [8, Ch. 8] and [35].

Here we focus on the case when  $A$  is of type (FPV), and  $B$  is maximally monotone such that

$$\text{star}(\text{dom } A) \cap \text{int dom } B \neq \emptyset.$$

(Implicitly this means that  $B$  is also of type (FPV).) In Theorem 3.3 we shall show that  $A + B$  is maximally monotone. As noted it seems possible that all maximally monotone operators are of type (FPV).

The remainder of this paper is organized as follows. In Section 2, we collect auxiliary results for future reference and for the reader’s convenience. In Section 3, our main result (Theorem 3.3) is presented. In Section 4, we then provide various corollaries and examples. We also pose several significant open questions on the sum problem. We leave the details of proof of Case 2 of Theorem 3.3 to Appendix 5.

## 2 Auxiliary Results

We first introduce one of Rockafellar’s results.

**Fact 2.1 (Rockafellar)** (See [23, Theorem 1] or [28, Theorem 27.1 and Theorem 27.3].) *Let  $A : X \rightrightarrows X^*$  be maximally monotone with  $\text{int dom } A \neq \emptyset$ . Then  $\text{int dom } A = \text{int } \overline{\text{dom } A}$  and  $\text{int dom } A$  and  $\overline{\text{dom } A}$  are both convex.*

The Fitzpatrick function defined below has proven to be an important tool in Monotone Operator Theory.

**Fact 2.2 (Fitzpatrick)** (See [17, Corollary 3.9].) *Let  $A : X \rightrightarrows X^*$  be monotone, and set*

$$(1) \quad F_A : X \times X^* \rightarrow ]-\infty, +\infty] : (x, x^*) \mapsto \sup_{(a, a^*) \in \text{gra } A} (\langle x, a^* \rangle + \langle a, x^* \rangle - \langle a, a^* \rangle),$$

the Fitzpatrick function associated with  $A$ . Suppose also  $A$  is maximally monotone. Then for every  $(x, x^*) \in X \times X^*$ , the inequality  $\langle x, x^* \rangle \leq F_A(x, x^*)$  is true, and the equality holds if and only if  $(x, x^*) \in \text{gra } A$ .

The next result is central to our arguments.

**Fact 2.3** (See [33, Theorem 3.4 and Corollary 5.6], or [28, Theorem 24.1(b)].) *Let  $A, B : X \rightrightarrows X^*$  be maximally monotone operators. Assume  $\bigcup_{\lambda > 0} \lambda [P_X(\text{dom } F_A) - P_X(\text{dom } F_B)]$  is a closed subspace. If*

$$F_{A+B} \geq \langle \cdot, \cdot \rangle \text{ on } X \times X^*,$$

*then  $A + B$  is maximally monotone.*

We next cite several results regarding operators of type (FPV).

**Fact 2.4 (Simons)** (See [28, Theorem 46.1].) *Let  $A : X \rightrightarrows X^*$  be a maximally monotone linear relation. Then  $A$  is of type (FPV).*

The following result presents a sufficient condition for a maximally monotone operator to be of type (FPV).

**Fact 2.5 (Simons and Verona-Verona)** (See [28, Theorem 44.1], [29] or [5].) *Let  $A : X \rightrightarrows X^*$  be maximally monotone. Suppose that for every closed convex subset  $C$  of  $X$  with  $\text{dom } A \cap \text{int } C \neq \emptyset$ , the operator  $A + N_C$  is maximally monotone. Then  $A$  is of type (FPV).*

**Fact 2.6** (See [2, Lemma 2.5].) *Let  $C$  be a nonempty closed convex subset of  $X$  such that  $\text{int } C \neq \emptyset$ . Let  $c_0 \in \text{int } C$  and suppose that  $z \in X \setminus C$ . Then there exists  $\lambda \in ]0, 1[$  such that  $\lambda c_0 + (1 - \lambda)z \in \text{bdry } C$ .*

**Fact 2.7 (Boundedness below)** (See [9, Fact 4.1].) *Let  $A : X \rightrightarrows X^*$  be monotone and  $x \in \text{int dom } A$ . Then there exist  $\delta > 0$  and  $M > 0$  such that  $x + \delta B_X \subseteq \text{dom } A$  and  $\sup_{a \in x + \delta B_X} \|Aa\| \leq M$ . Assume that  $(z, z^*)$  is monotonically related to  $\text{gra } A$ . Then*

$$\langle z - x, z^* \rangle \geq \delta \|z^*\| - (\|z - x\| + \delta)M.$$

**Fact 2.8 (Voisei and Zălinescu)** (See [36, Corollary 4].) *Let  $A, B : X \rightrightarrows X^*$  be maximally monotone. Assume that  ${}^{ic}(\text{dom } A) \neq \emptyset, {}^{ic}(\text{dom } B) \neq \emptyset$  and  $0 \in {}^{ic}[\text{dom } A - \text{dom } B]$ . Then  $A + B$  is maximally monotone.*

The proof of the next Lemma 2.9 follows closely the lines of that of [12, Lemma 2.10]. It generalizes both [12, Lemma 2.10] and [3, Lemma 2.10].

**Lemma 2.9** *Let  $A : X \rightrightarrows X^*$  be monotone, and let  $B : X \rightrightarrows X^*$  be a maximally monotone operator. Suppose that  $\text{star}(\text{dom } A) \cap \text{int dom } B \neq \emptyset$ . Suppose also that  $(z, z^*) \in X \times X^*$  with  $z \in \text{dom } A$  is monotonically related to  $\text{gra}(A + B)$ . Then  $z \in \text{dom } B$ .*

*Proof.* We can and do suppose that  $(0, 0) \in \text{gra } A \cap \text{gra } B$  and  $0 \in \text{star}(\text{dom } A) \cap \text{int dom } B$ . Suppose to the contrary that  $z \notin \text{dom } B$ . Then we have  $z \neq 0$ . We claim that

$$(2) \quad N_{[0,z]} + B \quad \text{is maximally monotone.}$$

Because  $z \neq 0$ , we have  $\frac{1}{2}z \in {}^{ic}(\text{dom } N_{[0,z]})$ . Clearly,  ${}^{ic}(\text{dom } B) \neq \emptyset$  and  $0 \in {}^{ic}[\text{dom } N_{[0,z]} - \text{dom } B]$ . By Fact 2.8,  $N_{[0,z]} + B$  is maximally monotone and hence (2) holds. Since  $(z, z^*) \notin \text{gra}(N_{[0,z]} + B)$ , there exist  $\lambda \in [0, 1]$  and  $x^*, y^* \in X^*$  such that  $(\lambda z, x^*) \in \text{gra } N_{[0,z]}$ ,  $(\lambda z, y^*) \in \text{gra } B$  and

$$(3) \quad \langle z - \lambda z, z^* - x^* - y^* \rangle < 0.$$

Now  $\lambda < 1$ , since  $(\lambda z, y^*) \in \text{gra } B$  and  $z \notin \text{dom } B$ , by (3),

$$(4) \quad \langle z, -x^* \rangle + \langle z, z^* - y^* \rangle = \langle z, z^* - x^* - y^* \rangle < 0.$$

Since  $(\lambda z, x^*) \in \text{gra } N_{[0,z]}$ , we have  $\langle z - \lambda z, x^* \rangle \leq 0$ . Then  $\langle z, -x^* \rangle \geq 0$ . Thus (4) implies that

$$(5) \quad \langle z, z^* - y^* \rangle < 0.$$

Since  $0 \in \text{star}(\text{dom } A)$  and  $z \in \text{dom } A$ ,  $\lambda z \in \text{dom } A$ . By the assumption on  $(z, z^*)$ , we have

$$\langle z - \lambda z, z^* - a^* - y^* \rangle \geq 0, \quad \forall a^* \in A(\lambda z).$$

Thence,  $\langle z, z^* - a^* - y^* \rangle \geq 0$  and hence

$$(6) \quad \langle z, z^* - y^* \rangle \geq \langle z, a^* \rangle, \quad \forall a^* \in A(\lambda z).$$

Next we show that

$$(7) \quad \langle z, a^* \rangle \geq 0, \quad \exists a^* \in A(\lambda z).$$

We consider two cases.

*Case 1:*  $\lambda = 0$ . Then take  $a^* = 0$  to see that (7) holds.

*Case 2:*  $\lambda \neq 0$ . Let  $a^* \in A(\lambda z)$ . Since  $(\lambda z, a^*) \in \text{gra } A$ ,  $\langle \lambda z, a^* \rangle = \langle \lambda z - 0, a^* - 0 \rangle \geq 0$  and hence  $\langle z, a^* \rangle \geq 0$ . Hence (7) holds.

Combining (6) and (7),

$$\langle z, z^* - y^* \rangle \geq 0, \quad \text{which contradicts (5).}$$

Hence  $z \in \text{dom } B$ . ■

The proof of Lemma 2.10 is modelled on that of [37, Proposition 3.1]. It is the first in a sequence of lemmas we give that will allow us to apply Fact 2.3.

**Lemma 2.10** *Let  $A : X \rightrightarrows X^*$  be monotone, and let  $B : X \rightrightarrows X^*$  be maximally monotone. Let  $(z, z^*) \in X \times X^*$ . Suppose  $x_0 \in \text{dom } A \cap \text{int dom } B$  and that there exists a sequence  $(a_n, a_n^*)_{n \in \mathbb{N}}$  in  $\text{gra } A \cap (\text{dom } B \times X^*)$  such that  $(a_n)_{n \in \mathbb{N}}$  converges to a point in  $[x_0, z[$ , while*

$$(8) \quad \langle z - a_n, a_n^* \rangle \longrightarrow +\infty.$$

Then  $F_{A+B}(z, z^*) = +\infty$ .

*Proof.* Since  $a_n \in \text{dom } B$  for every  $n \in \mathbb{N}$ , we may pick  $v_n^* \in B(a_n)$ . We again consider two cases.

*Case 1:*  $(v_n^*)_{n \in \mathbb{N}}$  is bounded.

Then we have

$$\begin{aligned} F_{A+B}(z, z^*) &\geq \sup_{\{n \in \mathbb{N}\}} [\langle a_n, z^* \rangle + \langle z - a_n, a_n^* \rangle + \langle z - a_n, v_n^* \rangle] \\ &\geq \sup_{\{n \in \mathbb{N}\}} [-\|a_n\| \cdot \|z^*\| + \langle z - a_n, a_n^* \rangle - \|z - a_n\| \cdot \|v_n^*\|] \\ &= +\infty \quad (\text{by (8) and the boundedness of } (v_n^*)_{n \in \mathbb{N}}). \end{aligned}$$

Hence  $F_{A+B}(z, z^*) = +\infty$ .

*Case 2:*  $(v_n^*)_{n \in \mathbb{N}}$  is unbounded.

By assumption, there exists  $0 \leq \lambda < 1$  such that

$$(9) \quad a_n \longrightarrow x_0 + \lambda(z - x_0).$$

We first show that

$$(10) \quad \limsup_{n \rightarrow \infty} \langle z - a_n, v_n^* \rangle = +\infty.$$

Since  $(v_n^*)_{n \in \mathbb{N}}$  is unbounded and, after passing to a subsequence if necessary, we may assume that  $\|v_n^*\| \neq 0, \forall n \in \mathbb{N}$  and that  $\|v_n^*\| \rightarrow +\infty$ . By  $x_0 \in \text{int dom } B$  and Fact 2.7, there exist  $\delta_0 > 0$  and  $K_0 > 0$  such that

$$(11) \quad \langle a_n - x_0, v_n^* \rangle \geq \delta_0 \|v_n^*\| - (\|a_n - x_0\| + \delta_0) K_0.$$

Then we have

$$(12) \quad \langle a_n - x_0, \frac{v_n^*}{\|v_n^*\|} \rangle \geq \delta_0 - \frac{(\|a_n - x_0\| + \delta_0) K_0}{\|v_n^*\|}, \quad \forall n \in \mathbb{N}.$$

By the Banach-Alaoglu Theorem (see [26, Theorem 3.15]), there exist a weak\* convergent subnet  $(\frac{v_\gamma^*}{\|v_\gamma^*\|})_{\gamma \in \Gamma}$  of  $(\frac{v_n^*}{\|v_n^*\|})_{n \in \mathbb{N}}$  such that

$$(13) \quad \frac{v_\gamma^*}{\|v_\gamma^*\|} \rightharpoonup_{w^*} v_\infty^* \in X^*.$$

Using (9) and taking the limit in (12) along the subnet, we obtain

$$(14) \quad \langle \lambda(z - x_0), v_\infty^* \rangle \geq \delta_0.$$

Hence  $\lambda$  is strictly positive and

$$(15) \quad \langle z - x_0, v_\infty^* \rangle \geq \frac{\delta_0}{\lambda} > 0.$$

Now assume contrary to (10) that there exists  $M > 0$  such that

$$\limsup_{n \rightarrow \infty} \langle z - a_n, v_n^* \rangle < M.$$

Then, for all  $n$  sufficiently large,

$$\langle z - a_n, v_n^* \rangle < M + 1,$$

and so

$$(16) \quad \langle z - a_n, \frac{v_n^*}{\|v_n^*\|} \rangle < \frac{M + 1}{\|v_n^*\|}.$$

Then by (9) and (13), taking the limit in (16) along the subnet again, we see that

$$(1 - \lambda)\langle z - x_0, v_\infty^* \rangle \leq 0.$$

Since  $\lambda < 1$ , we see  $\langle z - x_0, v_\infty^* \rangle \leq 0$  contradicting (15), and (10) holds. By (8) and (10),

$$F_{A+B}(z, z^*) \geq \sup_{n \in \mathbb{N}} [\langle a_n, z^* \rangle + \langle z - a_n, a_n \rangle + \langle z - a_n, v_n^* \rangle] = +\infty.$$

Hence

$$F_{A+B}(z, z^*) = +\infty,$$

as asserted. ■

We also need the following two lemmas.

**Lemma 2.11** *Let  $A : X \rightrightarrows X^*$  be monotone, and let  $B : X \rightrightarrows X^*$  be maximally monotone. Let  $(z, z^*) \in X \times X^*$ . Suppose that  $x_0 \in \text{dom } A \cap \text{int dom } B$  and that there exists a sequence  $(a_n)_{n \in \mathbb{N}}$  in  $\text{dom } A \cap \text{dom } B$  such that  $(a_n)_{n \in \mathbb{N}}$  converges to a point in  $[x_0, z[$ , and that*

$$(17) \quad a_n \in \text{bdry dom } B, \quad \forall n \in \mathbb{N}.$$

*Then  $F_{A+B}(z, z^*) = +\infty$ .*

*Proof.* Suppose to the contrary that

$$(18) \quad (z, z^*) \in \text{dom } F_{A+B}.$$

By the assumption, there exists  $0 \leq \lambda < 1$  such that

$$(19) \quad a_n \longrightarrow x_0 + \lambda(z - x_0).$$

By the Separation Theorem and Fact 2.1, there exists  $(y_n^*)_{n \in \mathbb{N}}$  in  $X^*$  such that  $\|y_n^*\| = 1$  and  $y_n^* \in N_{\overline{\text{dom } B}}(a_n)$ . Thus  $ky_n^* \in N_{\overline{\text{dom } B}}(a_n), \forall k > 0$ . Since  $x_0 \in \text{int dom } B$ , there exists  $\delta > 0$  such that  $x_0 + \delta B_X \subseteq \text{dom } B$ . Thus

$$\begin{aligned} \langle y_n^*, a_n \rangle &\geq \sup \langle y_n^*, x_0 + \delta B_X \rangle \geq \langle y_n^*, x_0 \rangle + \sup \langle y_n^*, \delta B_X \rangle = \langle y_n^*, x_0 \rangle + \delta \|y_n^*\| \\ &= \langle y_n^*, x_0 \rangle + \delta. \end{aligned}$$

Hence

$$(20) \quad \langle y_n^*, a_n - x_0 \rangle \geq \delta.$$

By the Banach-Alaoglu Theorem (see [26, Theorem 3.15]), there exists a weak\* convergent and bounded subnet  $(y_i^*)_{i \in O}$  such that

$$(21) \quad y_i^* \rightharpoonup_{w^*} y_\infty^* \in X^*.$$

Then (20) and (19) imply that

$$\langle y_\infty^*, \lambda(z - x_0) \rangle \geq \delta.$$

Thus, as before,  $\lambda > 0$  and

$$(22) \quad \langle y_\infty^*, z - x_0 \rangle \geq \frac{\delta}{\lambda} > 0.$$

Since  $B$  is maximally monotone,  $B = B + N_{\overline{\text{dom } B}}$ . As  $a_n \in \text{dom } A \cap \text{dom } B$ , we have

$$F_{A+B}(z, z^*) \geq \sup \left[ \langle z - a_n, A(a_n) \rangle + \langle z - a_n, B(a_n) + ky_n^* \rangle + \langle z^*, a_n \rangle \right], \quad \forall n \in \mathbb{N}, \forall k > 0.$$

Thus

$$\frac{F_{A+B}(z, z^*)}{k} \geq \sup \left[ \left\langle z - a_n, \frac{A(a_n)}{k} \right\rangle + \left\langle z - a_n, \frac{B(a_n)}{k} + y_n^* \right\rangle + \frac{\langle z^*, a_n \rangle}{k} \right], \quad \forall n \in \mathbb{N}, \forall k > 0.$$

Since  $(z, z^*) \in \text{dom } F_{A+B}$  by (18), on letting  $k \longrightarrow +\infty$  we obtain

$$0 \geq \langle z - a_n, y_n^* \rangle, \quad \forall n \in \mathbb{N}.$$

Combining with (21), (19) and taking the limit along the bounded subnet in the above inequality, we have

$$0 \geq \langle (1 - \lambda)(z - x_0), y_\infty^* \rangle.$$

Since  $\lambda < 1$ ,

$$\langle z - x_0, y_\infty^* \rangle \leq 0,$$

which contradicts (22).

Hence  $F_{A+B}(z, z^*) = +\infty$ . ■

**Lemma 2.12** *Let  $A : X \rightrightarrows X^*$  be of type (FPV). Suppose  $x_0 \in \text{dom } A$  but that  $z \notin \overline{\text{dom } A}$ . Then there is a sequence  $(a_n, a_n^*)_{n \in \mathbb{N}}$  in  $\text{gra } A$  so that  $(a_n)_{n \in \mathbb{N}}$  converges to a point in  $[x_0, z[$  and*

$$\langle z - a_n, a_n^* \rangle \longrightarrow +\infty.$$

*Proof.* Since  $z \notin \overline{\text{dom } A}$ ,  $z \neq x_0$ . Thence there exist  $\alpha > 0$  and  $y_0^* \in X^*$  such that  $\langle y_0^*, z - x_0 \rangle \geq \alpha$ . Set

$$U_n := [x_0, z] + \frac{1}{n}U_X, \quad \forall n \in \mathbb{N}.$$

Since  $x_0 \in \text{dom } A$ ,  $U_n \cap \text{dom } A \neq \emptyset$ . Now  $(z, ny_0^*) \notin \text{gra } A$  and  $z \in U_n$ . As  $A$  is of type (FPV), there exist  $(a_n, a_n^*)_{n \in \mathbb{N}}$  in  $\text{gra } A$  with  $a_n \in U_n$  such that

$$(23) \quad \langle z - a_n, a_n^* \rangle > \langle z - a_n, ny_0^* \rangle.$$

As  $a_n \in U_n$ ,  $(a_n)_{n \in \mathbb{N}}$  has a subsequence convergent to an element in  $[x_0, z]$ . We can assume that

$$(24) \quad a_n \longrightarrow x_0 + \lambda(z - x_0), \quad \text{where } 0 \leq \lambda \leq 1,$$

and since  $z \notin \overline{\text{dom } A}$ , we have  $\lambda < 1$ . Thus,  $x_0 + \lambda(z - x_0) \in [x_0, z[$ .

Thus by (24) and  $\langle z - x_0, y_0^* \rangle \geq \alpha > 0$ ,

$$\langle z - a_n, y_0^* \rangle \longrightarrow (1 - \lambda)\langle z - x_0, y_0^* \rangle \geq (1 - \lambda)\alpha > 0.$$

Hence there exists  $N_0 \in \mathbb{N}$  such that for every  $n \geq N_0$

$$(25) \quad \langle z - a_n, y_0^* \rangle \geq \frac{(1 - \lambda)\alpha}{2} > 0.$$

Appealing to (23), we have

$$\langle z - a_n, a_n^* \rangle > \frac{(1 - \lambda)\alpha}{2}n > 0, \quad \forall n \geq N_0,$$

and so  $\langle z - a_n, a_n^* \rangle \longrightarrow +\infty$ . This completes the proof. ■

### 3 Our main result

Before we come to our main result, we need the following two technical results which let us place points in the closures of the domains of  $A$  and  $B$ . The proof of Proposition 3.1 follows in part that of [37, Theorem 3.4].

**Proposition 3.1** *Let  $A : X \rightrightarrows X^*$  be of type (FPV), and let  $B : X \rightrightarrows X^*$  be maximally monotone. Suppose that  $\text{dom } A \cap \text{int dom } B \neq \emptyset$ . Let  $(z, z^*) \in X \times X^*$  with  $z \in \overline{\text{dom } B}$ . Then*

$$F_{A+B}(z, z^*) \geq \langle z, z^* \rangle.$$

*Proof.* Clearly,  $F_{A+B}(z, z^*) \geq \langle z, z^* \rangle$  if  $(z, z^*) \notin \text{dom } F_{A+B}$ . Now suppose that  $(z, z^*) \in \text{dom } F_{A+B}$ . We can suppose that  $0 \in \text{dom } A \cap \text{int dom } B$  and  $(0, 0) \in \text{gra } A \cap \text{gra } B$ . Next, we show that

$$(26) \quad F_{A+B}(tz, tz^*) \geq t^2 \langle z, z^* \rangle \quad \text{and} \quad tz \in \text{int dom } B, \quad \forall t \in ]0, 1[.$$

Fix  $t \in ]0, 1[$ . As  $0 \in \text{int dom } B$ ,  $z \in \overline{\text{dom } B}$ , Fact 2.1 and [40, Theorem 1.1.2(ii)] imply

$$(27) \quad tz \in \overline{\text{int dom } B},$$

and Fact 2.1 strengthens this to

$$(28) \quad tz \in \text{int dom } B.$$

We again consider two cases.

*Case 1:*  $tz \in \text{dom } A$ .

On selecting  $a^* \in A(tz), b^* \in B(tz)$ , the definition of the Fitzpatrick function (1) shows

$$F_{A+B}(tz, tz^*) \geq \langle tz^*, tz \rangle + \langle tz, a^* + b^* \rangle - \langle tz, a^* + b^* \rangle = \langle tz, tz^* \rangle.$$

Hence (26) holds.

*Case 2:*  $tz \notin \text{dom } A$ .

If  $\langle z, z^* \rangle \leq 0$ , then  $F_{A+B}(tz, tz^*) \geq 0 \geq \langle tz, tz^* \rangle$  because  $(0, 0) \in \text{gra } A \cap \text{gra } B$ . So we assume that

$$(29) \quad \langle z, z^* \rangle > 0.$$

We first show that

$$(30) \quad tz \in \overline{\text{dom } A}.$$

Set

$$U_n := [0, tz] + \frac{1}{n}U_X, \quad \forall n \in \mathbb{N}.$$

Since  $0 \in \text{dom } A$ ,  $U_n \cap \text{dom } A \neq \emptyset$ . Since  $(tz, nz^*) \notin \text{gra } A$  and  $tz \in U_n$ , while  $A$  is of type (FPV), there is  $(a_n, a_n^*)_{n \in \mathbb{N}}$  in  $\text{gra } A$  with  $a_n \in U_n$  such that

$$(31) \quad \langle tz, a_n^* \rangle > n \langle tz - a_n, z^* \rangle + \langle a_n, a_n^* \rangle.$$

As  $a_n \in U_n$ ,  $(a_n)_{n \in \mathbb{N}}$  has a subsequence convergent to an element in  $[0, tz]$ . We can assume that

$$(32) \quad a_n \longrightarrow \lambda z, \quad \text{where } 0 \leq \lambda \leq t.$$

As  $tz \in \text{int dom } B$  also  $\lambda z \in \text{int dom } B$ , and so appealing to Fact 2.7, there exist  $N \in \mathbb{N}$  and  $K > 0$  such that

$$(33) \quad a_n \in \text{int dom } B \quad \text{and} \quad \sup_{v^* \in B(a_n)} \|v^*\| \leq K, \quad \forall n \geq N.$$

We claim that

$$(34) \quad \lambda = t.$$

Suppose to the contrary that  $0 \leq \lambda < t$ . As  $(a_n, a_n^*) \in \text{gra } A$  and (33) holds, for every  $n \geq N$

$$\begin{aligned} & F_{A+B}(z, z^*) \\ & \geq \sup_{\{v^* \in B(a_n)\}} [\langle a_n, z^* \rangle + \langle z, a_n^* \rangle - \langle a_n, a_n^* \rangle + \langle z - a_n, v^* \rangle] \\ & \geq \sup_{\{v^* \in B(a_n)\}} [\langle a_n, z^* \rangle + \langle z, a_n^* \rangle - \langle a_n, a_n^* \rangle - K \|z - a_n\|] \\ & \geq \langle a_n, z^* \rangle + \langle z, a_n^* \rangle - \langle a_n, a_n^* \rangle - K \|z - a_n\| \\ & > \langle a_n, z^* \rangle + \frac{1}{t} n \langle tz - a_n, z^* \rangle + \frac{1}{t} \langle a_n, a_n^* \rangle - \langle a_n, a_n^* \rangle - K \|z - a_n\| \quad (\text{by (31)}) \\ & \geq \langle a_n, z^* \rangle + \frac{1}{t} n \langle tz - a_n, z^* \rangle - K \|z - a_n\| \quad (\text{since } \langle a_n, a_n^* \rangle \geq 0 \text{ by } (0, 0) \in \text{gra } A \text{ and } t \leq 1). \end{aligned}$$

Divide by  $n$  on both sides of the above inequality and take the limit with respect to  $n$ . Since (32) and  $F_{A+B}(z, z^*) < +\infty$ , we obtain

$$(1 - \frac{\lambda}{t}) \langle z, z^* \rangle = \langle z - \frac{\lambda}{t} z, z^* \rangle \leq 0.$$

Since  $0 \leq \lambda < t$ , we obtain  $\langle z, z^* \rangle \leq 0$ , which contradicts (29). Hence  $\lambda = t$  and by (32)  $tz \in \overline{\text{dom } A}$  so that (30) holds.

We next show that

$$(35) \quad F_{A+B}(tz, tz^*) \geq t^2 \langle z, z^* \rangle.$$

Set

$$H_n := tz + \frac{1}{n} U_X, \quad \forall n \in \mathbb{N}.$$

Note that  $H_n \cap \text{dom } A \neq \emptyset$ , since  $tz \in \overline{\text{dom } A} \setminus \text{dom } A$  by (30).

Because  $(tz, tz^*) \notin \text{gra } A$  and  $tz \in H_n$ , and  $A$  is of type (FPV), there exists  $(b_n, b_n^*)_{n \in \mathbb{N}}$  in  $\text{gra } A$  such that  $b_n \in H_n$  and

$$(36) \quad \langle tz, b_n^* \rangle + \langle b_n, tz^* \rangle - \langle b_n, b_n^* \rangle > t^2 \langle z, z^* \rangle, \quad \forall n \in \mathbb{N}.$$

As  $tz \in \text{int dom } B$  and  $b_n \rightarrow tz$ , by Fact 2.7, there exist  $N_1 \in \mathbb{N}$  and  $M > 0$  such that

$$(37) \quad b_n \in \text{int dom } B \quad \text{and} \quad \sup_{v^* \in B(b_n)} \|v^*\| \leq M, \quad \forall n \geq N_1.$$

We now compute

$$(38) \quad \begin{aligned} F_{A+B}(tz, tz^*) &\geq \sup_{\{c^* \in B(b_n)\}} [\langle b_n, tz^* \rangle + \langle tz, b_n^* \rangle - \langle b_n, b_n^* \rangle + \langle tz - b_n, c^* \rangle], \quad \forall n \geq N_1 \\ &\geq \sup_{\{c^* \in B(b_n)\}} [t^2 \langle z, z^* \rangle + \langle tz - b_n, c^* \rangle], \quad \forall n \geq N_1 \quad (\text{by (36)}) \\ &\geq \sup [t^2 \langle z, z^* \rangle - M \|tz - b_n\|], \quad \forall n \geq N_1 \quad (\text{by (37)}). \end{aligned}$$

Thus,

$$F_{A+B}(tz, tz^*) \geq t^2 \langle z, z^* \rangle$$

because  $b_n \rightarrow tz$ . Hence  $F_{A+B}(tz, tz^*) \geq t^2 \langle z, z^* \rangle$ . Thus (35) holds.

Combining the above cases, we see that (26) holds. Since  $(0, 0) \in \text{gra}(A + B)$  and  $A + B$  is monotone, we have  $F_{A+B}(0, 0) = \langle 0, 0 \rangle = 0$ . Since  $F_{A+B}$  is convex, (26) implies that

$$tF_{A+B}(z, z^*) = tF_{A+B}(z, z^*) + (1-t)F_{A+B}(0, 0) \geq F_{A+B}(tz, tz^*) \geq t^2 \langle z, z^* \rangle, \quad \forall t \in ]0, 1[.$$

Letting  $t \rightarrow 1^-$  in the above inequality, we obtain  $F_{A+B}(z, z^*) \geq \langle z, z^* \rangle$ . ■

We have one more block to put in place:

**Proposition 3.2** *Let  $A : X \rightrightarrows X^*$  be of type (FPV), and let  $B : X \rightrightarrows X^*$  be maximally monotone. Suppose  $\text{star}(\text{dom } A) \cap \text{int dom } B \neq \emptyset$ , and  $(z, z^*) \in \text{dom } F_{A+B}$ . Then  $z \in \overline{\text{dom } A}$ .*

*Proof.* We can and do suppose that  $0 \in \text{star}(\text{dom } A) \cap \text{int dom } B$  and  $(0, 0) \in \text{gra } A \cap \text{gra } B$ . As before, we suppose to the contrary that

$$(39) \quad z \notin \overline{\text{dom } A}.$$

Then  $z \neq 0$ . By the assumption that  $z \notin \overline{\text{dom } A}$ , Lemma 2.12 implies that there exist  $(a_n, a_n^*)_{n \in \mathbb{N}}$  in  $\text{gra } A$  and  $0 \leq \lambda < 1$  such that

$$(40) \quad \langle z - a_n, a_n^* \rangle \rightarrow +\infty \quad \text{and} \quad a_n \rightarrow \lambda z.$$

We yet again consider two cases.

*Case 1:* There exists a subsequence of  $(a_n)_{n \in \mathbb{N}}$  in  $\text{dom } B$ .

We can suppose that  $a_n \in \text{dom } B$  for every  $n \in \mathbb{N}$ . Thus by (40) and Lemma 2.10, we have  $F_{A+B}(z, z^*) = +\infty$ , which contradicts our original assumption that  $(z, z^*) \in \text{dom } F_{A+B}$ .

*Case 2:* There exists  $N_1 \in \mathbb{N}$  such that  $a_n \notin \text{dom } B$  for every  $n \geq N_1$ .

Now we can suppose that  $a_n \notin \text{dom } B$  for every  $n \in \mathbb{N}$ . Since  $a_n \notin \text{dom } B$ , Fact 2.1 and Fact 2.6 shows that there exists  $\lambda_n \in [0, 1]$  such that

$$(41) \quad \lambda_n a_n \in \text{bdry } \overline{\text{dom } B}.$$

By (40), we can suppose that

$$(42) \quad \lambda_n a_n \longrightarrow \lambda_\infty z.$$

Since  $0 \in \text{star}(\text{dom } A)$  and  $a_n \in \text{dom } A$ ,  $\lambda_n a_n \in \text{dom } A$ . Then (40) implies that

$$(43) \quad \lambda_\infty < 1.$$

We further split Case 2 into two subcases.

*Subcase 2.1:* There exists a subsequence of  $(\lambda_n a_n)_{n \in \mathbb{N}}$  in  $\text{dom } B$ . We may again suppose  $\lambda_n a_n \in \text{dom } B$  for every  $n \in \mathbb{N}$ . Since  $0 \in \text{star}(\text{dom } A)$  and  $a_n \in \text{dom } A$ ,  $\lambda_n a_n \in \text{dom } A$ . Then by (41) and (42), (43) and Lemma 2.11,  $F_{A+B}(z, z^*) = +\infty$ , which contradicts the hypothesis that  $(z, z^*) \in \text{dom } F_{A+B}$ .

*Subcase 2.2:* There exists  $N_2 \in \mathbb{N}$  such that  $\lambda_n a_n \notin \text{dom } B$  for every  $n \geq N_2$ . We can now assume that  $\lambda_n a_n \notin \text{dom } B$  for every  $n \in \mathbb{N}$ . Thus  $a_n \neq 0$  for every  $n \in \mathbb{N}$ . Since  $0 \in \text{int dom } B$ , (41) and (42) imply that  $0 < \lambda_\infty$  and then by (43)

$$(44) \quad 0 < \lambda_\infty < 1.$$

Since  $0 \in \text{int dom } B$ , (41) implies that  $\lambda_n > 0$  for every  $n \in \mathbb{N}$ . By (40),  $\|a_n - z\| \rightarrow 0$ . Then we can and do suppose that  $\|a_n - z\| \neq 0$  for every  $n \in \mathbb{N}$ . Fix  $n \in \mathbb{N}$ . Since  $0 \in \text{int dom } B$ , there exists  $0 < \rho_0 \leq 1$  such that  $\rho_0 B_X \subseteq \text{dom } B$ . As  $0 \in \text{star}(\text{dom } A)$  and  $a_n \in \text{dom } A$ ,  $\lambda_n a_n \in \text{dom } A$ . Set

$$(45) \quad b_n := \lambda_n a_n \quad \text{and take} \quad b_n^* \in A(\lambda_n a_n).$$

Next we show that there exists  $\varepsilon_n \in ]0, \frac{1}{n}[$  such that with  $H_n := (1 - \varepsilon_n)b_n + \varepsilon_n \rho_0 U_X$  and  $\tau_0 := \frac{1}{\lambda_n} \left[ 2\|z\| + 2\|a_n\| + 2 + (\|a_n\| + 1) \frac{2\lambda_n \|z - a_n\|}{\rho_0} \right]$ , we have

$$(46) \quad H_n \subseteq \text{dom } B \text{ and } \inf \|B(H_n)\| \geq n(1 + \tau_0 \|b_n^*\|), \text{ while } \varepsilon_n \max\{\|a_n\|, 1\} < \frac{1}{2} \|z - a_n\| \lambda_n.$$

For every  $s \in ]0, 1[$ , (41) and Fact 2.1 imply that  $(1 - s)b_n + s\rho_0 B_X \subseteq \overline{\text{dom } B}$ . By Fact 2.1 again,  $(1 - s)b_n + s\rho_0 U_X \subseteq \text{int } \overline{\text{dom } B} = \text{int dom } B$ .

Now we show the second assertion of (46). Let  $k \in \mathbb{N}$  and  $(s_k)_{k \in \mathbb{N}}$  be a positive sequence such that  $s_k \rightarrow 0$  when  $k \rightarrow \infty$ . It suffices to show

$$(47) \quad \liminf_{k \rightarrow \infty} \|B((1 - s_k)b_n + s_k \rho_0 U_X)\| = +\infty.$$

Suppose to the contrary there exist a sequence  $(c_k, c_k^*)_{k \in \mathbb{N}}$  in  $\text{gra } B \cap [((1 - s_k)b_n + s_k \rho_0 U_X) \times X^*]$  and  $L > 0$  such that  $\sup_{k \in \mathbb{N}} \|c_k^*\| \leq L$ . Then  $c_k \rightarrow b_n = \lambda_n a_n$ . By the Banach-Alaoglu Theorem (again see [26, Theorem 3.15]), there exist a weak\* convergent subnet,  $(c_\beta^*)_{\beta \in J}$  of  $(c_k^*)_{k \in \mathbb{N}}$  such that  $c_\beta^* \rightharpoonup^* c_\infty^* \in X^*$ . [9, Corollary 4.1] shows that  $(\lambda_n a_n, c_\infty^*) \in \text{gra } B$ , which contradicts our assumption that  $\lambda_n a_n \notin \text{dom } B$ . Hence (47) holds and so does (46).

Set  $t_n := \frac{\varepsilon_n \rho_0}{2\lambda_n \|z - a_n\|}$  and thus  $0 < t_n < \frac{1}{4}$ . Thus

$$(48) \quad t_n \lambda_n z + (1 - t_n)(1 - \varepsilon_n)b_n \in H_n.$$

Next we show there exists  $(\widetilde{a}_n, \widetilde{a}_n^*)_{n \in \mathbb{N}}$  in  $\text{gra } A \cap (H_n \times X^*)$  such that

$$(49) \quad \langle z - \widetilde{a}_n, \widetilde{a}_n^* \rangle \geq -\tau_0 \|b_n^*\|.$$

We consider two further subcases.

*Subcase 2.2a:*  $(t_n \lambda_n z + (1 - t_n)(1 - \varepsilon_n)b_n, (1 + t_n)b_n^*) \in \text{gra } A$ . Set  $(\widetilde{a}_n, \widetilde{a}_n^*) := (t_n \lambda_n z + (1 - t_n)(1 - \varepsilon_n)b_n, (1 + t_n)b_n^*)$ . Since  $(0, 0) \in \text{gra } A$ ,  $\langle b_n, b_n^* \rangle \geq 0$ .

Then we have

$$(50) \quad \begin{aligned} \langle t_n \lambda_n z - \widetilde{a}_n, \widetilde{a}_n^* \rangle &= \langle t_n \lambda_n z - t_n \lambda_n z - (1 - t_n)(1 - \varepsilon_n)b_n, (1 + t_n)b_n^* \rangle \\ &= \langle -(1 - t_n)(1 - \varepsilon_n)b_n, (1 + t_n)b_n^* \rangle = -\langle (1 - t_n^2)(1 - \varepsilon_n)b_n, b_n^* \rangle \geq -\langle b_n, b_n^* \rangle. \end{aligned}$$

On the other hand, (45) and the monotonicity of  $A$  imply that

$$\langle t_n \lambda_n z + (1 - t_n)(1 - \varepsilon_n)b_n - b_n, t_n b_n^* \rangle = \langle t_n \lambda_n z + (1 - t_n)(1 - \varepsilon_n)b_n - b_n, (1 + t_n)b_n^* - b_n^* \rangle \geq 0.$$

Thus

$$(51) \quad \langle t_n \lambda_n z - [1 - (1 - t_n)(1 - \varepsilon_n)]b_n, b_n^* \rangle \geq 0.$$

Since  $1 - (1 - t_n)(1 - \varepsilon_n) > 0$  and  $\langle b_n, b_n^* \rangle = \langle b_n - 0, b_n^* - 0 \rangle \geq 0$ , (51) implies that  $\langle t_n \lambda_n z, b_n^* \rangle \geq 0$  and thus

$$\langle z, b_n^* \rangle \geq 0.$$

Then by  $\widetilde{a}_n^* = (1 + t_n)b_n^*$  and  $t_n \lambda_n \leq 1$ , (50) implies that

$$\langle z - \widetilde{a}_n, \widetilde{a}_n^* \rangle \geq -\langle b_n, b_n^* \rangle \geq -\|b_n\| \cdot \|b_n^*\| \geq -\|a_n\| \cdot \|b_n^*\| \geq -\tau_0 \|b_n^*\|.$$

Hence (49) holds.

*Subcase 2.2b:*  $(t_n \lambda_n z + (1-t_n)(1-\varepsilon_n)b_n, (1+t_n)b_n^*) \notin \text{gra } A$ . By  $0 \in \text{star}(\text{dom } A)$  and  $a_n \in \text{dom } A$ , we have  $(1-\varepsilon_n)\lambda_n a_n \in \text{dom } A$ , hence  $\text{dom } A \cap H_n \neq \emptyset$ . Since  $t_n \lambda_n z + (1-t_n)(1-\varepsilon_n)b_n \in H_n$  by (48),  $(t_n \lambda_n z + (1-t_n)(1-\varepsilon_n)b_n, (1+t_n)b_n^*) \notin \text{gra } A$  and  $A$  is of type (FPV), there exists  $(\widetilde{a}_n, \widetilde{a}_n^*) \in \text{gra } A$  such that  $\widetilde{a}_n \in H_n$  and

$$\begin{aligned}
& \langle t_n \lambda_n z + (1-t_n)(1-\varepsilon_n)b_n - \widetilde{a}_n, \widetilde{a}_n^* - (1+t_n)b_n^* \rangle > 0 \\
& \Rightarrow \langle t_n \lambda_n z - [1 - (1-t_n)(1-\varepsilon_n)]\widetilde{a}_n + (1-t_n)(1-\varepsilon_n)(b_n - \widetilde{a}_n), \widetilde{a}_n^* - b_n^* \rangle \\
& \quad > \langle t_n \lambda_n z + (1-t_n)(1-\varepsilon_n)b_n - \widetilde{a}_n, t_n b_n^* \rangle \geq \langle t_n \lambda_n z - \widetilde{a}_n, t_n b_n^* \rangle \quad (\text{since } \langle b_n, b_n^* \rangle \geq 0) \\
& \Rightarrow \langle t_n \lambda_n z - [1 - (1-t_n)(1-\varepsilon_n)]\widetilde{a}_n, \widetilde{a}_n^* - b_n^* \rangle \\
& \quad > \langle (1-t_n)(1-\varepsilon_n)(b_n - \widetilde{a}_n), b_n^* - \widetilde{a}_n^* \rangle + \langle t_n \lambda_n z - \widetilde{a}_n, t_n b_n^* \rangle \\
& \Rightarrow \langle t_n \lambda_n z - [1 - (1-t_n)(1-\varepsilon_n)]\widetilde{a}_n, \widetilde{a}_n^* - b_n^* \rangle > \langle t_n \lambda_n z - \widetilde{a}_n, t_n b_n^* \rangle \\
& \Rightarrow \langle t_n \lambda_n z - [t_n + \varepsilon_n - t_n \varepsilon_n]\widetilde{a}_n, \widetilde{a}_n^* \rangle > \langle t_n \lambda_n z - \widetilde{a}_n, t_n b_n^* \rangle + \langle t_n \lambda_n z - [t_n + \varepsilon_n - t_n \varepsilon_n]\widetilde{a}_n, b_n^* \rangle.
\end{aligned}$$

Since  $\langle \widetilde{a}_n, \widetilde{a}_n^* \rangle = \langle \widetilde{a}_n - 0, \widetilde{a}_n^* - 0 \rangle \geq 0$  and  $t_n + \varepsilon_n - t_n \varepsilon_n \geq t_n \geq t_n \lambda_n$ ,  $\langle [t_n + \varepsilon_n - t_n \varepsilon_n]\widetilde{a}_n, \widetilde{a}_n^* \rangle \geq t_n \lambda_n \langle \widetilde{a}_n, \widetilde{a}_n^* \rangle$ . Thus

$$\begin{aligned}
& \langle t_n \lambda_n z - t_n \lambda_n \widetilde{a}_n, \widetilde{a}_n^* \rangle > \langle t_n \lambda_n z - \widetilde{a}_n, t_n b_n^* \rangle + \langle t_n \lambda_n z - [t_n + \varepsilon_n - t_n \varepsilon_n]\widetilde{a}_n, b_n^* \rangle \\
& \Rightarrow \langle \frac{t_n \lambda_n z - t_n \lambda_n \widetilde{a}_n}{\lambda_n t_n}, \widetilde{a}_n^* \rangle > \langle t_n \lambda_n z - \widetilde{a}_n, \frac{1}{\lambda_n} b_n^* \rangle + \langle \frac{t_n \lambda_n z - [t_n + \varepsilon_n - t_n \varepsilon_n]\widetilde{a}_n}{\lambda_n t_n}, b_n^* \rangle \\
& \Rightarrow \langle z - \widetilde{a}_n, \widetilde{a}_n^* \rangle > \langle t_n \lambda_n z - \widetilde{a}_n, \frac{1}{\lambda_n} b_n^* \rangle + \langle z - \left[1 + \frac{\varepsilon_n}{t_n} - \varepsilon_n\right] \frac{1}{\lambda_n} \widetilde{a}_n, b_n^* \rangle \\
& \Rightarrow \langle z - \widetilde{a}_n, \widetilde{a}_n^* \rangle > -\frac{1}{\lambda_n} \|b_n^*\| (\|z\| + \|a_n\| + 1) - \|b_n^*\| \left( \|z\| + \frac{1}{\lambda_n} (\|a_n\| + 1) \left(1 + \frac{2\lambda_n \|z - a_n\|}{\rho_0}\right) \right) \\
& \Rightarrow \langle z - \widetilde{a}_n, \widetilde{a}_n^* \rangle > -\|b_n^*\| \frac{1}{\lambda_n} \left[ 2\|z\| + 2\|a_n\| + 2 + (\|a_n\| + 1) \frac{2\lambda_n \|z - a_n\|}{\rho_0} \right] = -\tau_0 \|b_n^*\|.
\end{aligned}$$

Finally, combining all the subcases, we deduce that (49) holds.

Since  $\varepsilon_n < \frac{1}{n}$  and  $\widetilde{a}_n \in H_n$ , (42) shows that

$$(52) \quad \widetilde{a}_n \longrightarrow \lambda_\infty z.$$

Take  $w_n^* \in B(\widetilde{a}_n)$  by (46). Then by (46) again,

$$(53) \quad \|w_n^*\| \geq n(1 + \tau_0 \|b_n^*\|), \quad \forall n \in \mathbb{N}.$$

Then by (49), we have

$$\begin{aligned} -\tau_0 \|b_n^*\| + \left\langle z - \widetilde{a}_n, w_n^* \right\rangle + \left\langle z^*, \widetilde{a}_n \right\rangle &\leq \left\langle z - \widetilde{a}_n, \widetilde{a}_n^* \right\rangle + \left\langle z - \widetilde{a}_n, w_n^* \right\rangle + \left\langle z^*, \widetilde{a}_n \right\rangle \\ &\leq F_{A+B}(z, z^*) \end{aligned}$$

Thus

$$(54) \quad -\frac{\tau_0 \|b_n^*\|}{\|w_n^*\|} + \left\langle z - \widetilde{a}_n, \frac{w_n^*}{\|w_n^*\|} \right\rangle + \left\langle \frac{z^*}{\|w_n^*\|}, \widetilde{a}_n \right\rangle \leq \frac{F_{A+B}(z, z^*)}{\|w_n^*\|}.$$

By the Banach-Alaoglu Theorem (see [26, Theorem 3.15]), there exist a weak\* convergent subnet,  $(\frac{w_i^*}{\|w_i^*\|})_{i \in I}$  of  $\frac{w_n^*}{\|w_n^*\|}$  such that

$$(55) \quad \frac{w_i^*}{\|w_i^*\|} \xrightarrow{w^*} w_\infty^* \in X^*.$$

Combine (52), (53) and (55), by  $F_{A+B}(z, z^*) < +\infty$ , and take the limit along the subnet in (54) to obtain

$$\left\langle z - \lambda_\infty z, w_\infty^* \right\rangle \leq 0.$$

Then (44) shows that

$$(56) \quad \left\langle z, w_\infty^* \right\rangle \leq 0.$$

On the other hand, since  $0 \in \text{int dom } B$ , Fact 2.7 implies that there exists  $\rho_1 > 0$  and  $M > 0$  such that

$$\left\langle \widetilde{a}_n, w_n^* \right\rangle \geq \rho_1 \|w_n^*\| - (\|\widetilde{a}_n\| + \rho_1)M.$$

Thus

$$\left\langle \widetilde{a}_n, \frac{w_n^*}{\|w_n^*\|} \right\rangle \geq \rho_1 - \frac{(\|\widetilde{a}_n\| + \rho_1)M}{\|w_n^*\|}.$$

Use (52), (53) and (55), and take the limit along the subnet in the above inequality to obtain

$$\left\langle \lambda_\infty z, w_\infty^* \right\rangle \geq \rho_1.$$

Hence

$$\left\langle z, w_\infty^* \right\rangle \geq \frac{\rho_1}{\lambda_\infty} > 0,$$

which contradicts (56).

Combining all the above cases, we have arrived at  $z \in \overline{\text{dom } A}$ . ■

We are finally ready to prove our main result. The special case in which  $B$  is the normal cone operator of a nonempty closed convex set was first established by Voisei in [34].

**Theorem 3.3 ((FPV) Sum Theorem)** *Let  $A, B : X \rightrightarrows X^*$  be maximally monotone with  $\text{star}(\text{dom } A) \cap \text{int dom } B \neq \emptyset$ . Assume that  $A$  is of type (FPV). Then  $A+B$  is maximally monotone.*

*Proof.* After translating the graphs if necessary, we can and do assume that  $0 \in \text{star}(\text{dom } A) \cap \text{int dom } B$  and that  $(0, 0) \in \text{gra } A \cap \text{gra } B$ . By Fact 2.2,  $\text{dom } A \subseteq P_X(\text{dom } F_A)$  and  $\text{dom } B \subseteq P_X(\text{dom } F_B)$ . Hence,

$$(57) \quad \bigcup_{\lambda > 0} \lambda(P_X(\text{dom } F_A) - P_X(\text{dom } F_B)) = X.$$

Thus, by Fact 2.3, it suffices to show that

$$(58) \quad F_{A+B}(z, z^*) \geq \langle z, z^* \rangle, \quad \forall (z, z^*) \in X \times X^*.$$

Take  $(z, z^*) \in X \times X^*$ . Then

$$(59) \quad \begin{aligned} & F_{A+B}(z, z^*) \\ &= \sup_{\{x, x^*, y^*\}} [\langle x, z^* \rangle + \langle z - x, x^* \rangle + \langle z - x, y^* \rangle - \iota_{\text{gra } A}(x, x^*) - \iota_{\text{gra } B}(x, y^*)]. \end{aligned}$$

Suppose to the contrary that there exists  $\eta > 0$  such that

$$(60) \quad F_{A+B}(z, z^*) + \eta < \langle z, z^* \rangle,$$

so that

$$(61) \quad (z, z^*) \text{ is monotonically related to } \text{gra}(A+B).$$

Then by Proposition 3.1 and Proposition 3.2 ,

$$(62) \quad z \in \overline{\text{dom } A} \setminus \overline{\text{dom } B}.$$

Now by Lemma 2.9,

$$(63) \quad z \notin \text{dom } A.$$

Indeed, if  $z \in \text{dom } A$ , Lemma 2.9 and (61) show that  $z \in \text{dom } B$ . Thus,  $z \in \text{dom } A \cap \text{dom } B$  and hence  $F_{A+B}(z, z^*) \geq \langle z, z^* \rangle$  which contradicts (60). Thence we have established (63).

Thus (62) implies that there exists  $(a_n, a_n^*)_{n \in \mathbb{N}}$  in  $\text{gra } A$  such that

$$(64) \quad a_n \longrightarrow z.$$

By (62),  $a_n \notin \overline{\text{dom } B}$  for all but finitely many terms  $a_n$ . We can suppose that  $a_n \notin \overline{\text{dom } B}$  for all  $n \in \mathbb{N}$ . Fact 2.1 and Fact 2.6 show that there exists  $\lambda_n \in ]0, 1[$  such that

$$(65) \quad \lambda_n a_n \in \text{bdry } \overline{\text{dom } B}.$$

By (64), we can assume that

$$(66) \quad \lambda_n \longrightarrow \lambda_\infty \in [0, 1] \quad \text{and thus} \quad \lambda_n a_n \longrightarrow \lambda_\infty z.$$

Then by (65) and (62)

$$(67) \quad \lambda_\infty < 1.$$

We consider two cases.

*Case 1:* There exists a subsequence of  $(\lambda_n a_n)_{n \in \mathbb{N}}$  in  $\text{dom } B$ .

We can suppose that  $\lambda_n a_n \in \text{dom } B$  for every  $n \in \mathbb{N}$ . Since  $0 \in \text{star}(\text{dom } A)$  and  $a_n \in \text{dom } A$ ,  $\lambda_n a_n \in \text{dom } A$ . Then by (65), (66), (67) and Lemma 2.11,  $F_{A+B}(z, z^*) = +\infty$ , which contradicts (60) that  $(z, z^*) \in \text{dom } F_{A+B}$ .

*Case 2:* There exists  $N \in \mathbb{N}$  such that  $\lambda_n a_n \notin \text{dom } B$  for every  $n \geq N$ .

We can suppose that  $\lambda_n a_n \notin \text{dom } B$  for every  $n \in \mathbb{N}$ . Thus  $a_n \neq 0$  for every  $n \in \mathbb{N}$ . Following the pattern of Subcase 2.2 in the proof of Proposition 3.2<sup>1</sup>, we obtain a contradiction.

Combing all the above cases, we have  $F_{A+B}(z, z^*) \geq \langle z, z^* \rangle$  for all  $(z, z^*) \in X \times X^*$ . Hence  $A + B$  is maximally monotone.  $\blacksquare$

**Remark 3.4** In Case 2 in the proof of Theorem 3.3 (see Appendix 5 below), we use Lemma 2.9 to deduce that  $\|a_n - z\| \neq 0$ . Without the help of Lemma 2.9, we may still can obtain (77) as follows. For the case of  $a_n = z$ , consider whether  $((1 - \varepsilon_n)b_n, 0) = ((1 - \varepsilon_n)\lambda_n z, 0) \in H_n \times X^*$  is in  $\text{gra } A$  or not. We can deduce that there exists  $(\widetilde{a}_n, \widetilde{a}_n^*)_{n \in \mathbb{N}}$  in  $\text{gra } A \cap (H_n \times X^*)$  such that

$$\langle z - \widetilde{a}_n, \widetilde{a}_n^* \rangle \geq 0.$$

Hence (77) holds, and the proof of Theorem 3.3 can be achieved without Lemma 2.9.  $\diamond$

## 4 Examples and Consequences

We start by illustrating that the starshaped hypothesis catches operators whose domain may be non-convex and have no algebraic interior.

**Example 4.1 (Operators with starshaped domains)** We illustrate that there are many choices of maximally monotone operator  $A$  of type (FPV) with non-convex domain such that  ${}^{ic} \text{dom } A = \text{int dom } A = \emptyset$  and  $\text{star}(\text{dom } A) \neq \emptyset$ . Let  $f : \mathbb{R}^2 \rightarrow ]-\infty, +\infty]$  be defined by

$$(x, y) \mapsto \begin{cases} \max\{1 - \sqrt{x}, |y|\} & \text{if } x \geq 0; \\ +\infty, & \text{otherwise.} \end{cases}$$

---

<sup>1</sup>We banish the details to Appendix 5 to spare the readers.

Consider an infinite dimensional Banach space  $X$  containing a nonempty closed and convex set  $C$  such that  ${}^{ic}C = \emptyset$ . It is not known whether all spaces have this property but all separable or reflexive spaces certainly do [8]. Define  $A : (\mathbb{R}^2 \times X) \rightrightarrows (\mathbb{R}^2 \times X^*)$  by

$$(v, w) \rightrightarrows \left( \partial f(v), \partial \iota_C(w) \right) = \partial F(v, w),$$

where  $F := f \oplus \iota_C$ . Define  $\|\cdot\|$  on  $\mathbb{R}^2 \times X$  by  $\|(v, w)\| := \|v\| + \|w\|$ .

Then  $f$  is proper convex and lowers semicontinuous and so, therefore, is  $F$ . Indeed, [22, Example before Theorem 23.5, page 218] shows that  $\text{dom } \partial f$  is not convex and consequently  $\text{dom } A$  is not convex. (Many other candidates for  $f$  are given in [8, Chapter 7].) Clearly,  $A = \partial F$  is maximally monotone. Let  $w_0 \in C$  and  $v_0 = (2, 0)$ . Consider  $(v_0, w_0) \in \mathbb{R}^2 \times X$ . Since  $v_0 = (2, 0) \in \text{int } \text{dom } \partial f$ ,  $v_0 \in \text{star}(\text{dom } \partial f)$  since  $\text{dom } f$  is convex. Thus  $(v_0, w_0) \in \text{star}(\text{dom } A)$ . Since  ${}^{ic}C = \emptyset$  and so  $\text{int } C = \emptyset$ , it follows that  ${}^{ic} \text{dom } A = \text{int } \text{dom } A = \emptyset$ . [28, Theorem 48.4(d)] shows that  $A = \partial F$  is of type (FPV).  $\diamond$

The next example gives all the details of how to associate the support points of a convex set to a subgradient. In [18], [13] and [8, Exercise 8.4.1, page 401] the construction is used to build empty subgradients in various Fréchet spaces and incomplete normed spaces.

**Example 4.2 (Support points)** Suppose that  $X$  is separable. We can always find a compact convex set  $C \subseteq X$  such that  $\text{span } C \neq X$  and  $\overline{\text{span } C} = X$  [8]. Take  $x_0 \notin \text{span } C$ . Define  $f : X \rightarrow ]-\infty, +\infty]$  by

$$(68) \quad f(x) := \min\{t \in \mathbb{R} \mid x + tx_0 \in C\}, \quad \forall x \in X.$$

By direct computation  $f$  is proper lower semicontinuous and convex, see [18]. By the definition of  $f$ ,  $\text{dom } f = C + \mathbb{R}x_0$ . Let  $t \in \mathbb{R}$  and  $c \in C$ . We shall establish that

$$(69) \quad \partial f(tx_0 + c) = \begin{cases} N_C(c) \cap \{y^* \in X^* \mid \langle y^*, x_0 \rangle = -1\}, & \text{if } c \in \text{supp } C; \\ \emptyset, & \text{otherwise.} \end{cases}$$

Thence, also  $\text{dom } \partial f = \mathbb{R}x_0 + \text{supp } C$ .

First we show that the implication

$$(70) \quad tx_0 + c = sx_0 + d, \quad \text{where } t, s \in \mathbb{R}, c, d \in C \quad \Rightarrow \quad t = s \quad \text{and} \quad c = d$$

holds. Let  $t, s \in \mathbb{R}$  and  $c, d \in C$ . We have  $(t - s)x_0 = d - c \in \text{span } C - \text{span } C = \text{span } C$ . Since  $x_0 \notin \text{span } C$ ,  $t = s$  and then  $c = d$ . Hence we obtain (70).

By (70), we have

$$(71) \quad f(tx_0 + c) = -t, \quad \forall t \in \mathbb{R}, \forall c \in C.$$

We next show that (69) holds.

Since  $\text{dom } f = C + \mathbb{R}x_0$ , by (71), we have

$$\begin{aligned}
& x^* \in \partial f(tx_0 + c) \\
& \Leftrightarrow \langle x^*, sx_0 + d - (tx_0 + c) \rangle \leq f(sx_0 + d) - f(tx_0 + c) = -s + t, \quad \forall s \in \mathbb{R}, \forall d \in C \\
& \Leftrightarrow \langle x^*, (s-t)x_0 + (d-c) \rangle \leq -s + t, \quad \forall s \in \mathbb{R}, \forall d \in C \\
& \Leftrightarrow \langle x^*, sx_0 + (d-c) \rangle \leq -s, \quad \forall s \in \mathbb{R}, \forall d \in C \\
& \Leftrightarrow \langle x^*, sx_0 \rangle \leq -s \quad \text{and} \quad \langle x^*, d-c \rangle \leq 0, \quad \forall s \in \mathbb{R}, \forall d \in C \\
& \Leftrightarrow \langle x^*, sx_0 \rangle \leq -s \quad \text{and} \quad x^* \in N_C(c), x^* \neq 0, \quad \forall s \in \mathbb{R} \\
& \Leftrightarrow \langle x^*, x_0 \rangle = -1, \quad x^* \in N_C(c) \quad \text{and} \quad c \in \text{supp } C.
\end{aligned}$$

Hence (69) holds.

As a concrete example of  $C$  consider, for  $1 \leq p < \infty$ , any order interval  $C := \{x \in \ell^p(\mathbb{N}) : \alpha \leq x \leq \beta\}$  where  $\alpha < \beta \in \ell^p(\mathbb{N})$ . The example extends to all weakly compactly generated (WCG) spaces [8] with a weakly compact convex set in the role of  $C$ .  $\diamond$

We gave the last example in part as it allows one to better understand what the domain of a maximally monotone operator with empty interior can look like. While the star may be empty, it has been recently proven [32], see also [16], that for a closed convex function  $f$  the domain of  $\partial f$  is always pathwise and locally pathwise connected.

An immediate corollary of Theorem 3.3 is the following which generalizes [37, Corollary 3.9].

**Corollary 4.3 (Convex domain)** *Let  $A, B : X \rightrightarrows X^*$  be maximally monotone with  $\text{dom } A \cap \text{int dom } B \neq \emptyset$ . Assume that  $A$  is of type (FPV) with convex domain. Then  $A + B$  is maximally monotone.*

An only slightly less immediate corollary is given next.

**Corollary 4.4 (Nonempty interior)** (See [5, Theorem 9(i)] or Fact 2.8.) *Let  $A, B : X \rightrightarrows X^*$  be maximally monotone with  $\text{int dom } A \cap \text{int dom } B \neq \emptyset$ . Then  $A + B$  is maximally monotone.*

*Proof.* By the assumption, there exists  $x_0 \in \text{int dom } A \cap \text{int dom } B$ . We first show that  $A$  is of type (FPV). Let  $C$  be a nonempty closed convex subset of  $X$ , and suppose that  $\text{dom } A \cap \text{int } C \neq \emptyset$ . Let  $x_1 \in \text{dom } A \cap \text{int } C$ . Fact 2.1 and [40, Theorem 1.1.2(ii)] imply that  $[x_0, x_1[ \subseteq \text{int dom } A = \text{int dom } A$ . Since  $x_1 \in \text{int } C$ , there exists  $0 < \delta < 1$  such that  $x_1 + \delta(x_0 - x_1) \in \text{int dom } A \cap \text{int } C$ . Then  $N_C + A$  is maximally monotone by Corollary 4.3 and [28, Theorem 48.4(d)]. Hence by Fact 2.5,  $A$  is of type (FPV), see also [5].

Since  $x_0 \in \text{int dom } A$ , Fact 2.1 and [40, Theorem 1.1.2(ii)] imply that  $x_0 \in \text{star}(\text{dom } A)$  and hence we have  $x_0 \in \text{star}(\text{dom } A) \cap \text{int dom } B$ . Then by Theorem 3.3, we deduce that  $A + B$  is maximally monotone. ■

**Corollary 4.5 (Linear relation)** (See [12, Theorem 3.1].) *Let  $A : X \rightrightarrows X^*$  be a maximally monotone linear relation, and let  $B : X \rightrightarrows X^*$  be maximally monotone. Suppose that  $\text{dom } A \cap \text{int dom } B \neq \emptyset$ . Then  $A + B$  is maximally monotone.*

*Proof.* Apply Fact 2.4 and Corollary 4.3 directly. ■

The proof of our final Corollary 4.6 is adapted from that of [37, Corollary 2.10] and [12, Corollary 3.3]. Moreover, it generalizes both [37, Corollary 2.10] and [12, Corollary 3.3].

**Corollary 4.6 (FPV property of the sum)** *Let  $A, B : X \rightrightarrows X^*$  be maximally monotone with  $\text{dom } A \cap \text{int dom } B \neq \emptyset$ . Assume that  $A$  is of type (FPV) with convex domain, as holds if  $A$  is type (FPV) with closed domain. Then  $A + B$  is of type (FPV).*

*Proof.* By Corollary 4.3,  $A + B$  is maximally monotone. Let  $C$  be a nonempty closed convex subset of  $X$ , and suppose that  $\text{dom}(A + B) \cap \text{int } C \neq \emptyset$ . Let  $x_1 \in \text{dom } A \cap \text{int dom } B$  and  $x_2 \in \text{dom}(A + B) \cap \text{int } C$ . Then  $x_1, x_2 \in \text{dom } A$ ,  $x_1 \in \text{int dom } B$  and  $x_2 \in \text{dom } B \cap \text{int } C$ . Hence  $\lambda x_1 + (1 - \lambda)x_2 \in \text{int dom } B$  for every  $\lambda \in ]0, 1]$  by Fact 2.1 and [40, Theorem 1.1.2(ii)] and so there exists  $\delta \in ]0, 1]$  such that  $\lambda x_1 + (1 - \lambda)x_2 \in \text{int } C$  for every  $\lambda \in [0, \delta]$ .

Thus,  $\delta x_1 + (1 - \delta)x_2 \in \text{dom } A \cap \text{int dom } B \cap \text{int } C$ . By Corollary 4.4,  $B + N_C$  is maximally monotone. Then, by Corollary 4.3 (applied  $A$  and  $B + N_C$  to  $A$  and  $B$ ),  $A + B + N_C = A + (B + N_C)$  is maximally monotone. By Fact 2.5,  $A + B$  is of type (FPV).

The final conclusion holds since as we have noted  $\text{dom } A$  has a convex closure. ■

We have been unable to relax the convexity hypothesis in Corollary 4.6.

We finish by listing some related interesting, at least to the current authors, questions regarding the sum problem.

**Open Problem 4.7** *Let  $A : X \rightrightarrows X^*$  be maximally monotone with convex domain. Is  $A$  necessarily of type (FPV)?*

Let us recall a problem posed by S. Simons in [27, Problem 41.2]

**Open Problem 4.8** *Let  $A : X \rightrightarrows X^*$  be of type (FPV), let  $C$  be a nonempty closed convex subset of  $X$ , and suppose that  $\text{dom } A \cap \text{int } C \neq \emptyset$ . Is  $A + N_C$  necessarily maximally monotone?*

More generally, can we relax or indeed entirely drop the starshaped hypothesis on  $\text{dom } A$  in Theorem 3.3?

**Open Problem 4.9** Let  $A, B : X \rightrightarrows X^*$  be maximally monotone with  $\text{dom } A \cap \text{int dom } B \neq \emptyset$ . Assume that  $A$  is of type (FPV). Is  $A + B$  necessarily maximally monotone?

If all maximally monotone operators are type (FPV) this is no easier than the full sum problem. Can the results of [32] help here?

## Acknowledgment

Jonathan Borwein and Liangjin Yao were partially supported by various Australian Research Council grants.

## References

- [1] H.H. Bauschke and P.L. Combettes, *Convex Analysis and Monotone Operator Theory in Hilbert Spaces*, Springer-Verlag, 2011.
- [2] H.H. Bauschke, X. Wang, and L. Yao, “An answer to S. Simons’ question on the maximal monotonicity of the sum of a maximal monotone linear operator and a normal cone operator”, *Set-Valued and Variational Analysis*, vol. 17, pp. 195–201, 2009.
- [3] H.H. Bauschke, X. Wang, and L. Yao, “On the maximal monotonicity of the sum of a maximal monotone linear relation and the subdifferential operator of a sublinear function”, *Proceedings of the Haifa Workshop on Optimization Theory and Related Topics. Contemp. Math., Amer. Math. Soc., Providence, RI*, vol. 568, pp. 19–26, 2012.
- [4] J.M. Borwein, “Maximal monotonicity via convex analysis”, *Journal of Convex Analysis*, vol. 13, pp. 561–586, 2006.
- [5] J.M. Borwein, “Maximality of sums of two maximal monotone operators in general Banach space”, *Proceedings of the American Mathematical Society*, vol. 135, pp. 3917–3924, 2007.
- [6] J.M. Borwein, “Fifty years of maximal monotonicity”, *Optimization Letters*, vol. 4, pp. 473–490, 2010.
- [7] J.M. Borwein and A.S. Lewis, *Convex Analysis and Nonsmooth Optimization*, Second expanded edition, Springer, 2005.
- [8] J.M. Borwein and J. Vanderwerff, *Convex Functions: Constructions, Characterizations and Counterexamples*, Encyclopedia of Mathematics and its Applications, **109**, Cambridge University Press, 2010.
- [9] J.M. Borwein and L. Yao, “Structure theory for maximally monotone operators with points of continuity”, *Journal of Optimization Theory and Applications*, vol 156, pp. 1–24, 2013 (Invited paper).

- [10] J.M. Borwein and L. Yao, “Some results on the convexity of the closure of the domain of a maximally monotone operator”, *Optimization Letters*, in press;  
<http://dx.doi.org/10.1007/s11590-012-0564-7>.
- [11] J.M. Borwein and L. Yao, “Recent progress on Monotone Operator Theory”, *Infinite Products of Operators and Their Applications*, Contemporary Mathematics, in press;  
<http://arxiv.org/abs/1210.3401v2>.
- [12] J.M. Borwein and L. Yao, “Maximality of the sum of a maximally monotone linear relation and a maximally monotone operator”, *Set-Valued and Variational Analysis*, in press;  
<http://arxiv.org/abs/1212.4266v1>.
- [13] A. Brøndsted and R.T. Rockafellar, “On the subdifferentiability of convex functions”, *Proceedings of the American Mathematical Society*, vol. 16, pp. 605–611, 1995.
- [14] R.S. Burachik and A.N. Iusem, *Set-Valued Mappings and Enlargements of Monotone Operators*, Springer-Verlag, 2008.
- [15] D. Butnariu and A.N. Iusem, *Totally Convex Functions for Fixed Points Computation and Infinite Dimensional Optimization*, Kluwer Academic Publishers, 2000.
- [16] C. De Bernardi and L. Veselý, “On support points and support functionals of convex sets,” *Israel J. Math.* vol. 171, pp. 15-27, 2009.
- [17] S. Fitzpatrick, “Representing monotone operators by convex functions”, in *Workshop/Miniconference on Functional Analysis and Optimization (Canberra 1988)*, Proceedings of the Centre for Mathematical Analysis, Australian National University, vol. 20, Canberra, Australia, pp. 59–65, 1988.
- [18] R. B. Holmes, *Geometric Functional Analysis and its Applications*, Springer-Verlag, New York, 1975.
- [19] M. Marques Alves and B.F. Svaiter, “A new qualification condition for the maximality of the sum of maximal monotone operators in general Banach spaces”, *Journal of Convex Analysis*, vol. 19, pp. 575–589, 2012.
- [20] R.R. Phelps, *Convex Functions, Monotone Operators and Differentiability*, 2nd Edition, Springer-Verlag, 1993.
- [21] R.R. Phelps and S. Simons, “Unbounded linear monotone operators on nonreflexive Banach spaces”, *Journal of Nonlinear and Convex Analysis*, vol. 5, pp. 303–328, 1998.
- [22] R.T. Rockafellar, *Convex Analysis*, Princeton Univ. Press, Princeton, 1970.
- [23] R.T. Rockafellar, “Local boundedness of nonlinear, monotone operators”, *Michigan Mathematical Journal*, vol. 16, pp. 397–407, 1969.
- [24] R.T. Rockafellar, “On the maximality of sums of nonlinear monotone operators”, *Transactions of the American Mathematical Society*, vol. 149, pp. 75–88, 1970.

- [25] R.T. Rockafellar and R.J-B Wets, *Variational Analysis*, 3rd Printing, Springer-Verlag, 2009.
- [26] R. Rudin, *Functional Analysis*, Second Edition, McGraw-Hill, 1991.
- [27] S. Simons, *Minimax and Monotonicity*, Springer-Verlag, 1998.
- [28] S. Simons, *From Hahn-Banach to Monotonicity*, Springer-Verlag, 2008.
- [29] A. Verona and M.E. Verona, “Regular maximal monotone operators”, *Set-Valued Analysis*, vol. 6, pp. 303–312, 1998.
- [30] A. Verona and M.E. Verona, “Regular maximal monotone operators and the sum theorem”, *Journal of Convex Analysis*, vol. 7, pp. 115–128, 2000.
- [31] A. Verona and M.E. Verona, “On the regularity of maximal monotone operators and related results”; <http://arxiv.org/abs/1212.1968v3>, December 2012.
- [32] L. Vesely “A parametric smooth variational principle and support properties of convex sets and functions”, *Journal of Mathematical Analysis and Applications*, vol. 350, pp. 550–561, 2009.
- [33] M.D. Voisei, “The sum and chain rules for maximal monotone operators”, *Set-Valued and Variational Analysis*, vol. 16, pp. 461–476, 2008.
- [34] M.D. Voisei, “A Sum Theorem for (FPV) operators and normal cones ”, *Journal of Mathematical Analysis and Applications*, vol. 371, pp. 661–664, 2010.
- [35] M.D. Voisei and C. Zălinescu, “Strongly-representable monotone operators”, *Journal of Convex Analysis* , vol. 16, pp. 1011–1033, 2009.
- [36] M.D. Voisei and C. Zălinescu, “Maximal monotonicity criteria for the composition and the sum under weak interiority conditions”, *Mathematical Programming (Series B)*, vol. 123, pp. 265–283, 2010.
- [37] L. Yao, “The sum of a maximal monotone operator of type (FPV) and a maximal monotone operator with full domain is maximally monotone”, *Nonlinear Anal.*, vol. 74, pp. 6144–6152, 2011.
- [38] L. Yao, “The sum of a maximally monotone linear relation and the subdifferential of a proper lower semicontinuous convex function is maximally monotone”, *Set-Valued and Variational Analysis*, vol. 20, pp. 155–167, 2012.
- [39] L. Yao, *On Monotone Linear Relations and the Sum Problem in Banach Spaces*, Ph.D. thesis, University of British Columbia (Okanagan), 2011; <http://hdl.handle.net/2429/39970>.
- [40] C. Zălinescu, *Convex Analysis in General Vector Spaces*, World Scientific Publishing, 2002.
- [41] E. Zeidler, *Nonlinear Functional Analysis and its Applications II/A: Linear Monotone Operators*, Springer-Verlag, 1990.
- [42] E. Zeidler, *Nonlinear Functional Analysis and its Applications II/B: Nonlinear Monotone Operators*, Springer-Verlag, 1990.

## 5 Appendix

### Proof of Case 2 in the proof of Theorem 3.3.

*Proof. Case 2:* There exists  $N \in \mathbb{N}$  such that  $\lambda_n a_n \notin \text{dom } B$  for every  $n \geq N$ .

We can and do suppose that  $\lambda_n a_n \notin \text{dom } B$  for every  $n \in \mathbb{N}$ . Thus  $a_n \neq 0$  for every  $n \in \mathbb{N}$ .

Since  $0 \in \text{int dom } B$ , (66) and (65) imply that  $0 < \lambda_\infty$  and hence by (67)

$$(72) \quad 0 < \lambda_\infty < 1.$$

By (63),  $\|a_n - z\| \neq 0$  for every  $n \in \mathbb{N}$ .

Fix  $n \in \mathbb{N}$ . Since  $0 \in \text{int dom } B$ , there exists  $0 < \rho_0 \leq 1$  such that  $\rho_0 B_X \subseteq \text{dom } B$ . Since  $0 \in \text{star}(\text{dom } A)$  and  $a_n \in \text{dim } A$ ,  $\lambda_n a_n \in \text{dom } A$ . Set

$$(73) \quad b_n := \lambda_n a_n \quad \text{and take} \quad b_n^* \in A(\lambda_n a_n).$$

Next we show that there exists  $\varepsilon_n \in ]0, \frac{1}{n}[$  such that

$$(74) \quad H_n \subseteq \text{dom } B \quad \text{and} \quad \inf \|B(H_n)\| \geq n(1 + \tau_0 \|b_n^*\|), \quad \varepsilon_n \max\{\|a_n\|, 1\} < \frac{1}{2} \|z - a_n\| \lambda_n.$$

where  $H_n := (1 - \varepsilon_n)b_n + \varepsilon_n \rho_0 U_X$  and  $\tau_0 := \frac{1}{\lambda_n} \left[ 2\|z\| + 2\|a_n\| + 2 + (\|a_n\| + 1) \frac{2\lambda_n \|z - a_n\|}{\rho_0} \right]$ .

For every  $\varepsilon \in ]0, 1[$ , by (65) and Fact 2.1,  $(1 - \varepsilon)b_n + \varepsilon \rho_0 B_X \subseteq \overline{\text{dom } B}$ . By Fact 2.1 again,  $(1 - \varepsilon)b_n + \varepsilon \rho_0 U_X \subseteq \text{int dom } B = \text{int dom } B$ .

Now we show the second part of (74). Let  $k \in \mathbb{N}$  and  $(s_k)_{k \in \mathbb{N}}$  be a positive sequence such that  $s_k \rightarrow 0$  when  $k \rightarrow \infty$ . It suffices to show

$$(75) \quad \liminf_{k \rightarrow \infty} \|B((1 - s_k)b_n + s_k \rho_0 U_X)\| = +\infty.$$

Suppose to the contrary there exist a sequence  $(c_k, c_k^*)_{k \in \mathbb{N}}$  in  $\text{gra } B \cap [((1 - s_k)b_n + s_k \rho_0 U_X) \times X^*]$  and  $L > 0$  such that  $\sup_{k \in \mathbb{N}} \|c_k^*\| \leq L$ . Then  $c_k \rightarrow b_n = \lambda_n a_n$ . By the Banach-Alaoglu Theorem (see [26, Theorem 3.15]), there exist a weak\* convergent subnet,  $(c_\beta^*)_{\beta \in J}$  of  $(c_k^*)_{k \in \mathbb{N}}$  such that  $c_\beta^* \xrightarrow{w^*} c_\infty^* \in X^*$ . [9, Corollary 4.1] shows that  $(\lambda_n a_n, c_\infty^*) \in \text{gra } B$ , which contradicts our assumption that  $\lambda_n a_n \notin \text{dom } B$ .

Hence (75) holds and so does (74).

Set  $t_n := \frac{\varepsilon_n \rho_0}{2\lambda_n \|z - a_n\|}$  and thus  $0 < t_n < \frac{1}{4}$ . Thus

$$(76) \quad t_n \lambda_n z + (1 - t_n)(1 - \varepsilon_n)b_n \in H_n.$$

Next we show there exists  $(\widetilde{a}_n, \widetilde{a}_n^*)_{n \in \mathbb{N}}$  in  $\text{gra } A \cap (H_n \times X^*)$  such that

$$(77) \quad \langle z - \widetilde{a}_n, \widetilde{a}_n^* \rangle \geq -\tau_0 \|b_n^*\|.$$

We consider two subcases.

*Subcase 2.1:*  $(t_n \lambda_n z + (1 - t_n)(1 - \varepsilon_n) b_n, (1 + t_n) b_n^*) \in \text{gra } A$ .

Then set  $(\widetilde{a}_n, \widetilde{a}_n^*) := (t_n \lambda_n z + (1 - t_n)(1 - \varepsilon_n) b_n, (1 + t_n) b_n^*)$ . Since  $(0, 0) \in \text{gra } A$ ,  $\langle b_n, b_n^* \rangle \geq 0$ . Then we have

$$(78) \quad \begin{aligned} & \langle t_n \lambda_n z - \widetilde{a}_n, \widetilde{a}_n^* \rangle = \langle t_n \lambda_n z - t_n \lambda_n z - (1 - t_n)(1 - \varepsilon_n) b_n, (1 + t_n) b_n^* \rangle \\ & = \langle -(1 - t_n)(1 - \varepsilon_n) b_n, (1 + t_n) b_n^* \rangle = -\langle (1 - t_n^2)(1 - \varepsilon_n) b_n, b_n^* \rangle \geq -\langle b_n, b_n^* \rangle. \end{aligned}$$

On the other hand, (73) and the monotonicity of  $A$  imply that

$$\langle t_n \lambda_n z + (1 - t_n)(1 - \varepsilon_n) b_n - b_n, t_n b_n^* \rangle = \langle t_n \lambda_n z + (1 - t_n)(1 - \varepsilon_n) b_n - b_n, (1 + t_n) b_n^* - b_n^* \rangle \geq 0$$

Thus

$$(79) \quad \langle t_n \lambda_n z - [1 - (1 - t_n)(1 - \varepsilon_n)] b_n, b_n^* \rangle \geq 0.$$

Since  $1 - (1 - t_n)(1 - \varepsilon_n) > 0$  and  $\langle b_n, b_n^* \rangle = \langle b_n - 0, b_n^* - 0 \rangle \geq 0$ , (79) implies that  $\langle t_n \lambda_n z, b_n^* \rangle \geq 0$  and thus

$$\langle z, b_n^* \rangle \geq 0.$$

Then by  $\widetilde{a}_n^* = (1 + t_n) b_n^*$  and  $t_n \lambda_n \leq 1$ , (78) implies that

$$\langle z - \widetilde{a}_n, \widetilde{a}_n^* \rangle \geq -\langle b_n, b_n^* \rangle \geq -\|b_n\| \cdot \|b_n^*\| \geq -\|a_n\| \cdot \|b_n^*\| \geq -\tau_0 \|b_n^*\|.$$

Hence (77) holds.

*Subcase 2.2:*  $(t_n \lambda_n z + (1 - t_n)(1 - \varepsilon_n) b_n, (1 + t_n) b_n^*) \notin \text{gra } A$ .

Since  $0 \in \text{star}(\text{dom } A)$  and  $a_n \in \text{dom } A$ , we have  $(1 - \varepsilon_n) \lambda_n a_n \in \text{dom } A$ , hence  $\text{dom } A \cap H_n \neq \emptyset$ . Since  $t_n \lambda_n z + (1 - t_n)(1 - \varepsilon_n) b_n \in H_n$  by (76),  $(t_n \lambda_n z + (1 - t_n)(1 - \varepsilon_n) b_n, (1 + t_n) b_n^*) \notin \text{gra } A$  and  $A$  is of type (FPV), there exists  $(\widetilde{a}_n, \widetilde{a}_n^*) \in \text{gra } A$  such that  $\widetilde{a}_n \in H_n$  and

$$\begin{aligned} & \langle t_n \lambda_n z + (1 - t_n)(1 - \varepsilon_n) b_n - \widetilde{a}_n, \widetilde{a}_n^* - (1 + t_n) b_n^* \rangle > 0 \\ & \Rightarrow \langle t_n \lambda_n z - [1 - (1 - t_n)(1 - \varepsilon_n)] \widetilde{a}_n + (1 - t_n)(1 - \varepsilon_n)(b_n - \widetilde{a}_n), \widetilde{a}_n^* - b_n^* \rangle \\ & > \langle t_n \lambda_n z + (1 - t_n)(1 - \varepsilon_n) b_n - \widetilde{a}_n, t_n b_n^* \rangle \geq \langle t_n \lambda_n z - \widetilde{a}_n, t_n b_n^* \rangle \quad (\text{since } \langle b_n, b_n^* \rangle \geq 0) \\ & \Rightarrow \langle t_n \lambda_n z - [1 - (1 - t_n)(1 - \varepsilon_n)] \widetilde{a}_n, \widetilde{a}_n^* - b_n^* \rangle \\ & > \langle (1 - t_n)(1 - \varepsilon_n)(b_n - \widetilde{a}_n), b_n^* - \widetilde{a}_n^* \rangle + \langle t_n \lambda_n z - \widetilde{a}_n, t_n b_n^* \rangle \\ & \Rightarrow \langle t_n \lambda_n z - [1 - (1 - t_n)(1 - \varepsilon_n)] \widetilde{a}_n, \widetilde{a}_n^* - b_n^* \rangle > \langle t_n \lambda_n z - \widetilde{a}_n, t_n b_n^* \rangle \end{aligned}$$

$$\Rightarrow \langle t_n \lambda_n z - [t_n + \varepsilon_n - t_n \varepsilon_n] \widetilde{a}_n, \widetilde{a}_n^* \rangle > \langle t_n \lambda_n z - \widetilde{a}_n, t_n b_n^* \rangle + \langle t_n \lambda_n z - [t_n + \varepsilon_n - t_n \varepsilon_n] \widetilde{a}_n, b_n^* \rangle.$$

Since  $\langle \widetilde{a}_n, \widetilde{a}_n^* \rangle = \langle \widetilde{a}_n - 0, \widetilde{a}_n^* - 0 \rangle \geq 0$  and  $t_n + \varepsilon_n - t_n \varepsilon_n \geq t_n \geq t_n \lambda_n$ ,  $\langle [t_n + \varepsilon_n - t_n \varepsilon_n] \widetilde{a}_n, \widetilde{a}_n^* \rangle \geq t_n \lambda_n \langle \widetilde{a}_n, \widetilde{a}_n^* \rangle$ . Thus

$$\begin{aligned} & \langle t_n \lambda_n z - t_n \lambda_n \widetilde{a}_n, \widetilde{a}_n^* \rangle > \langle t_n \lambda_n z - \widetilde{a}_n, t_n b_n^* \rangle + \langle t_n \lambda_n z - [t_n + \varepsilon_n - t_n \varepsilon_n] \widetilde{a}_n, b_n^* \rangle \\ & \Rightarrow \langle \frac{t_n \lambda_n z - t_n \lambda_n \widetilde{a}_n}{\lambda_n t_n}, \widetilde{a}_n^* \rangle > \langle t_n \lambda_n z - \widetilde{a}_n, \frac{1}{\lambda_n} b_n^* \rangle + \langle \frac{t_n \lambda_n z - [t_n + \varepsilon_n - t_n \varepsilon_n] \widetilde{a}_n}{\lambda_n t_n}, b_n^* \rangle \\ & \Rightarrow \langle z - \widetilde{a}_n, \widetilde{a}_n^* \rangle > \langle t_n \lambda_n z - \widetilde{a}_n, \frac{1}{\lambda_n} b_n^* \rangle + \langle z - \left[1 + \frac{\varepsilon_n}{t_n} - \varepsilon_n\right] \frac{1}{\lambda_n} \widetilde{a}_n, b_n^* \rangle \\ & \Rightarrow \langle z - \widetilde{a}_n, \widetilde{a}_n^* \rangle > -\frac{1}{\lambda_n} \|b_n^*\| (\|z\| + \|a_n\| + 1) - \|b_n^*\| \left( \|z\| + \frac{1}{\lambda_n} (\|a_n\| + 1) \left(1 + \frac{2\lambda_n \|z - a_n\|}{\rho_0}\right) \right) \\ & \Rightarrow \langle z - \widetilde{a}_n, \widetilde{a}_n^* \rangle > -\|b_n^*\| \frac{1}{\lambda_n} \left[ 2\|z\| + 2\|a_n\| + 2 + (\|a_n\| + 1) \frac{2\lambda_n \|z - a_n\|}{\rho_0} \right] = -\tau_0 \|b_n^*\|. \end{aligned}$$

Hence combining all the subcases, we have (77) holds.

Since  $\varepsilon_n < \frac{1}{n}$  and  $\widetilde{a}_n \in H_n$ , (66) shows that

$$(80) \quad \widetilde{a}_n \longrightarrow \lambda_\infty z.$$

Take  $w_n^* \in B(\widetilde{a}_n)$  by (74). Then by (74) again,

$$(81) \quad \|w_n^*\| \geq n(1 + \tau_0 \|b_n^*\|), \quad \forall n \in \mathbb{N}.$$

Then by (77), we have

$$\begin{aligned} -\tau_0 \|b_n^*\| + \langle z - \widetilde{a}_n, w_n^* \rangle + \langle z^*, \widetilde{a}_n \rangle & \leq \langle z - \widetilde{a}_n, \widetilde{a}_n^* \rangle + \langle z - \widetilde{a}_n, w_n^* \rangle + \langle z^*, \widetilde{a}_n \rangle \\ & \leq F_{A+B}(z, z^*) \end{aligned}$$

Thus

$$(82) \quad -\frac{\tau_0 \|b_n^*\|}{\|w_n^*\|} + \left\langle z - \widetilde{a}_n, \frac{w_n^*}{\|w_n^*\|} \right\rangle + \left\langle \frac{z^*}{\|w_n^*\|}, \widetilde{a}_n \right\rangle \leq \frac{F_{A+B}(z, z^*)}{\|w_n^*\|}.$$

By the Banach-Alaoglu Theorem (see [26, Theorem 3.15]), there exist a weak\* convergent subnet,  $(\frac{w_i^*}{\|w_i^*\|})_{i \in I}$  of  $\frac{w_n^*}{\|w_n^*\|}$  such that

$$(83) \quad \frac{w_i^*}{\|w_i^*\|} \rightharpoonup_{w^*} w_\infty^* \in X^*.$$

Combining (80), (81) and (83), by  $F_{A+B}(z, z^*) < +\infty$ , take the limit along the subnet in (82) to obtain

$$\left\langle z - \lambda_\infty z, w_\infty^* \right\rangle \leq 0.$$

Then (72) shows that

$$(84) \quad \langle z, w_\infty^* \rangle \leq 0.$$

On the other hand, since  $0 \in \text{int dom } B$ , Fact 2.7 implies that there exists  $\rho_1 > 0$  and  $M > 0$  such that

$$\langle \widetilde{a}_n, w_n^* \rangle \geq \rho_1 \|w_n^*\| - (\|\widetilde{a}_n\| + \rho_1)M.$$

Thus

$$\left\langle \widetilde{a}_n, \frac{w_n^*}{\|w_n^*\|} \right\rangle \geq \rho_1 - \frac{(\|\widetilde{a}_n\| + \rho_1)M}{\|w_n^*\|}.$$

Combining (80), (81) and (83), take the limit along the subnet in the above inequality to obtain

$$\langle \lambda_\infty z, w_\infty^* \rangle \geq \rho_1.$$

Hence

$$\langle z, w_\infty^* \rangle \geq \frac{\rho_1}{\lambda_\infty} > 0,$$

which contradicts (84). ■