

# A Sinc that Sank

David Borwein\*, Jonathan M. Borwein<sup>†</sup> and Armin Straub<sup>‡</sup>

CARMA Analysis Seminar, September 6, 2011

## Abstract

We resolve and further study a sinc integral evaluation, first posed in THE AMERICAN MATHEMATICAL MONTHLY in [1967, p. 1015], which was solved in [1968, p. 914] and withdrawn in [1970, p. 657].

After a short introduction to the problem and its history, we give a general evaluation which we make entirely explicit in the case of the product of three sinc functions.

Finally, we exhibit some general structure of the integrals in question.

**Key words:** sinc function, conditionally convergent integral, numeric-symbolic computation.

- The associated preprint is available at <http://carma.newcastle.edu.au/jon/sink.pdf>.


---

\*Department of Mathematics, University of Western Ontario, London ON Canada.

<sup>†</sup>Corresponding author, CARMA, University of Newcastle, NSW 2308 Australia and KAU, Jeddah, SA. Email: [jonathan.borwein@newcastle.edu.au](mailto:jonathan.borwein@newcastle.edu.au).

<sup>‡</sup>Tulane University, USA. Email: [astraub@tulane.edu](mailto:astraub@tulane.edu)

# Some More Scenes from a Scientist's Life ...



*"This could be the discovery of the century. Depending, of course, on how far down it goes."*

# 1 Introduction and background

In [1967, #5529, p. 1015] D. Mitrinovic asked in this MONTHLY for an evaluation of

$$I_n := \int_{-\infty}^{\infty} \prod_{j=1}^n \frac{\sin k_j(x - a_j)}{x - a_j} dx \quad (1)$$

for real numbers  $a_j, k_j$  with  $1 \leq j \leq n$ . We shall write  $I_n \left( \begin{smallmatrix} a_1, \dots, a_n \\ k_1, \dots, k_n \end{smallmatrix} \right)$  when we wish to emphasize the dependence on the parameters.

The next year a solution [1968, #5529, p. 914] was published in the form of

$$I_n = \pi \prod_{j=2}^n \frac{\sin k_j(a_1 - a_j)}{a_1 - a_j}. \quad (2)$$

This solution as Klamkin pointed out in [1970, p. 657] can not be correct, since it is not symmetric in the parameters while  $I_n$  is. Indeed  $k_1 = 0$  forces  $I_n = 0$ . The proof given relies on formally correct Fourier analysis; but there are missing constraints on the  $k_j$  variables which have the effect that it is seldom right for more than two variables. Indeed, as shown then by Djokvič and Glasser [7] — who were both working in Waterloo at the time — the evaluation (2) holds true under the restriction  $k_1 \geq k_2 + k_3 + \dots + k_n$  when all of the  $k_j$  are positive.

However, no simple general fix appeared possible — and indeed for  $n > 2$  the issue is somewhat complex — and the problem while recorded several times in later MONTHLY lists of unsolved problems appears (from a JSTOR hunt<sup>1</sup>) to have disappeared without trace in the later 1980's.

The precise issues regarding evaluation of sinc integrals are described in detail in [3] or [4, Chapter 2] along with some remarkable consequences [2, 3, 4]. In the two-variable case the 1968 solution is essentially correct: we do obtain

$$I_2 = \pi \frac{\sin(k_1 \wedge k_2)(a_1 - a_2)}{a_1 - a_2} \quad (3)$$

---

<sup>1</sup>A search through all MONTHLY volumes, suggests that the solutions were never published and indeed for some years the original problem reappeared on lists of unsolved MONTHLY problems before apparently disappearing from view. Such a JSTOR search is not totally convincing since there is no complete indexing of problems and their status.

for  $a_1 \neq a_2$  as will be made explicit below. Some of the delicacy is a consequence of the fact that the classical sinc evaluation —  $\text{sinc } x := \frac{\sin x}{x}$  — given next is only conditionally true [3]. We have

$$\int_{-\infty}^{\infty} \frac{\sin kx}{x} dx = \pi \operatorname{sgn}(k), \quad (4)$$

where  $\operatorname{sgn}(0) = 0$ ,  $\operatorname{sgn}(k) = 1$  for  $k > 0$  and  $\operatorname{sgn}(k) = -1$  for  $k < 0$ .

In (4) the integral is absolutely divergent and is best interpreted as a Cauchy-Riemann limit.

Thus, the evaluation of (1) yields  $I_1 = \pi \operatorname{sgn}(k_1)$  which has a discontinuity at  $k_1 = 0$ . For  $n \geq 2$ , however,  $I_n$  is an absolutely convergent integral which is (jointly) continuous as a function of all  $k_j$  and all  $a_j$ . This follows from Lebesgue's dominated convergence theorem since the absolute value of the integrand is less than  $\prod_{j=1}^n |k_j| + 1$  for all  $x$  and less than  $2/x^2$  for all sufficiently large  $|x|$ .

We finish the introduction by observing that the oscillatory structure of the integrals, see Figure 1, means that their evaluation both numerically and symbolically calls for a significant amount of care.




Figure 1: Integrand in (1) with  $\mathbf{a} = (-3, -2, -1, 0, 1, 2)$ ,  $\mathbf{k} = (1, 2, 3, 4, 5, 6)$

We also note the continuing fascination with similar sinc integrals [8]. Indeed, [3] was triggered by the same problem described by Morrison in this MONTHLY [10]. This led also to a lovely MONTHLY article on random series [11].

Finally, Feeman's recent book on medical imaging [5] chose to begin with the example given at the beginning of Section 4.

## 2 Evaluation of $I_n$

Without loss of generality we assume that all  $k_j$  are strictly positive. In this section we shall only consider the case when all the  $a_j$  are distinct. As illustrated in Sections 3.2 and 3.3 the special cases can be treated by taking limits. We begin with the classical and simple partial fraction expression

$$\prod_{j=1}^n \frac{1}{x - a_j} = \sum_{j=1}^n \frac{1}{x - a_j} \prod_{i \neq j} \frac{1}{a_j - a_i} \quad (5)$$

valid when the  $a_j$  are distinct. Applying (5) to the integral  $I_n$  we then have:

$$\begin{aligned} I_n &= \sum_{j=1}^n \int_{-\infty}^{\infty} \frac{\sin(k_j(x - a_j))}{x - a_j} \prod_{i \neq j} \frac{\sin(k_i(x - a_i))}{a_j - a_i} dx \\ &= \sum_{j=1}^n \int_{-\infty}^{\infty} \frac{\sin(k_j x)}{x} \prod_{i \neq j} \frac{\sin(k_i(x + (a_j - a_i)))}{a_j - a_i} dx \end{aligned} \quad (6)$$

We pause and illustrate the general approach in the case of  $n = 2$  variables.

**Example 2.1** (Two variables). We apply (6) to write

$$\begin{aligned} I_2 &= \int_{-\infty}^{\infty} \frac{\sin k_1 x}{x} \frac{\sin k_2(x + a_1 - a_2)}{a_2 - a_1} dx + \int_{-\infty}^{\infty} \frac{\sin k_2 x}{x} \frac{\sin k_1(x + a_2 - a_1)}{a_1 - a_2} dx \\ &= \frac{\sin k_2(a_1 - a_2)}{a_1 - a_2} \int_{-\infty}^{\infty} \frac{\sin k_1 x}{x} \cos(k_2 x) dx + \frac{\sin k_1(a_2 - a_1)}{a_2 - a_1} \int_{-\infty}^{\infty} \frac{\sin k_2 x}{x} \cos(k_1 x) dx \end{aligned}$$

where for the second equation we have used the addition formula for the sine and noticed that the sine terms (being odd) integrate to zero. Finally, we either appeal to [3, Theorem 3] or express

$$\begin{aligned} \int_{-\infty}^{\infty} \frac{\sin k_1 x}{x} \cos(k_2 x) dx &= \frac{1}{2} \int_{-\infty}^{\infty} \frac{\sin(k_1 + k_2)x}{x} dx + \frac{1}{2} \int_{-\infty}^{\infty} \frac{\sin(k_1 - k_2)x}{x} dx, \\ \int_{-\infty}^{\infty} \frac{\sin k_2 x}{x} \cos(k_1 x) dx &= \frac{1}{2} \int_{-\infty}^{\infty} \frac{\sin(k_1 + k_2)x}{x} dx - \frac{1}{2} \int_{-\infty}^{\infty} \frac{\sin(k_1 - k_2)x}{x} dx, \end{aligned}$$

and appeal twice to (4) to obtain the final elegant cancellation

$$I_2 = \pi \frac{\sin((k_1 \wedge k_2)(a_1 - a_2))}{a_1 - a_2} \quad (7)$$

valid for  $a_1 \neq a_2$ . Here  $a \wedge b := \min\{a, b\}$ . We observe that the result remains true for  $a_1 = a_2$ , in which case the right-hand side of (7) attains the limiting value  $k_1 \wedge k_2$ .  $\diamond$

Let us observe that after the first step in Example 2.1 — independent of the exact final formula — the integrals to be obtained have lost their dependence on the  $a_j$ . This is what we exploit more generally. Proceeding as in Example 2.1 and applying the addition formula to (6) we write:

$$I_n = \sum_{j=1}^n \sum_{A,B} C_{j,A,B} \int_{-\infty}^{\infty} \prod_{i \in A \cup \{j\}} \sin(k_i x) \prod_{i \in B} \cos(k_i x) \frac{dx}{x} \quad (8)$$

where the sum is over all sets  $A$  and  $B$  partitioning  $\{1, 2, \dots, j-1, j+1, \dots, n\}$ , and

$$C_{j,A,B} := \prod_{i \in A} \frac{\cos(k_i(a_j - a_i))}{a_j - a_i} \prod_{i \in B} \frac{\sin(k_i(a_j - a_i))}{a_j - a_i}. \quad (9)$$

Notice that we may assume the cardinality  $|A|$  of  $A$  to be even since the integral in (8) vanishes if  $|A|$  is odd.

To further treat (8) we write the products of sines and cosines in terms of sums of single trigonometric functions. The general formulae are made explicit next.

**Proposition 2.2** (Cosine Product).

$$\prod_{j=1}^n \cos(x_j) = 2^{-n} \sum_{\varepsilon \in \{-1, 1\}^n} \cos\left(\sum_{j=1}^n \varepsilon_j x_j\right). \quad (10)$$

*Proof.* The formula follows inductively from the trigonometric identity  $2 \cos(a) \cos(b) = \cos(a+b) + \cos(a-b)$ .  $\square$

Observe that by taking derivatives with respect to some of the  $x_j$  in (10) we obtain similar formulae for general products of sines and cosines.

**Corollary 2.3** (Sine and Cosine Product).

$$\prod_{j=1}^n \sin(x_j) \prod_{j=n+1}^{n+m} \cos(x_j) = 2^{-n} \sum_{\varepsilon \in \{-1,1\}^{n+m}} \left( \prod_{j=1}^n \varepsilon_j \right) \cos \left( \sum_{j=1}^{n+m} \varepsilon_j x_j - \frac{n\pi}{2} \right). \quad (11)$$

It follows that, for even  $|A|$ ,

$$\begin{aligned} \alpha_{j,A,B} &:= \int_{-\infty}^{\infty} \prod_{i \in A \cup \{j\}} \sin(k_i x) \prod_{i \in B} \cos(k_i x) \frac{dx}{x} \\ &= (-1)^{|A|/2} \int_{-\infty}^{\infty} \frac{1}{2^n} \sum_{\varepsilon \in \{-1,1\}^n} \left( \prod_{i \in A \cup \{j\}} \varepsilon_i \right) \sin \left( \sum_{i=1}^n \varepsilon_i k_i x \right) \frac{dx}{x} \\ &= \pi (-1)^{|A|/2} \frac{1}{2^n} \sum_{\varepsilon \in \{-1,1\}^n} \left( \prod_{i \in A \cup \{j\}} \varepsilon_i \right) \operatorname{sgn} \left( \sum_{i=1}^n \varepsilon_i k_i \right). \end{aligned} \quad (12)$$

Then on combining (12) with (8) we obtain the following general evaluation:

**Theorem 2.4** (General Evaluation). *We have*

$$I_n = \sum_{j=1}^n \sum_{A,B} \alpha_{j,A,B} C_{j,A,B} \quad (13)$$

where the inner summation is over disjoint sets  $A, B$  such that  $|A|$  is even and  $A \cup B = \{1, 2, \dots, j-1, j+1, \dots, n\}$ .

The trigonometric products

$$C_{j,A,B} = \prod_{i \in A} \frac{\cos(k_i(a_j - a_i))}{a_j - a_i} \prod_{i \in B} \frac{\sin(k_i(a_j - a_i))}{a_j - a_i}$$

are as in (9) and  $\alpha_{j,A,B}$  is given by (12).

Note that in dimension  $n \geq 2$ , there are  $n2^{n-2}$  elements  $C_{j,A,B}$  which may or may not be distinct.

**Remark 2.5.** Note that, just like for the defining integral for  $I_n$ , it is apparent that the terms  $C_{j,A,B}$  and hence the evaluation of  $I_n$  given in (13) only depend on the parameters  $a_j$  up to a common shift. In particular, setting  $b_j = a_j - a_n$  for  $j = 1, \dots, n-1$  the evaluation in (13) can be written as a symmetric function in the  $n-1$  variables  $b_j$ .  $\diamond$

As an immediate consequence of Theorem 2.4 we have:

**Corollary 2.6** (Simplest Case). *Assume, without loss, that  $k_1, k_2, \dots, k_n > 0$ . Suppose that there is an index  $\ell$  such that*

$$k_\ell > \frac{1}{2} \sum k_i.$$

*In that case, the original solution to the MONTHLY problem is valid; that is,*

$$I_n = \int_{-\infty}^{\infty} \prod_{i=1}^n \frac{\sin(k_i(x - a_i))}{x - a_i} dx = \pi \prod_{i \neq \ell} \frac{\sin(k_i(a_\ell - a_i))}{a_\ell - a_i}.$$

This result was independently obtained by Djokvič and Glasser [7].

*Proof.* In this case,

$$\operatorname{sgn} \left( \sum_{i=1}^n \varepsilon_i k_i \right) = \varepsilon_\ell$$

for all values of the  $\varepsilon_i$ . The claim now follows from Theorem 2.4. More precisely, if there is some index  $k \neq \ell$  such that  $k \in A$  or  $j = k$  then  $\alpha_{j,A,B} = 0$ . This is because the term in (12) contributed by  $\varepsilon \in \{-1, 1\}^n$  has opposite sign than the term contributed by  $\varepsilon'$ , where  $\varepsilon'$  is obtained from  $\varepsilon$  by flipping the sign of  $\varepsilon_k$ . It remains to observe that  $\alpha_{\ell, \emptyset, B} = \pi$ .  $\square$

## 2.1 Alternative evaluation of $I_n$

In 1970 Dragomir Ž. Djokvič sent in a solution to the MONTHLY after the original solution was withdrawn [7]. He used the following identity involving the principal value (PV) of the integral

$$(\text{PV}) \int_{-\infty}^{\infty} \frac{e^{itx}}{x - a_j} dx = \lim_{\delta \rightarrow 0^+} \left\{ \int_{-\infty}^{a_j - \delta} + \int_{a_j + \delta}^{\infty} \right\} \frac{e^{itx}}{x - a_j} dx = \pi i \operatorname{sgn}(t) e^{ita_j} \quad (14)$$

where  $t$  is real. Note that setting  $a_j = 0$  and taking the imaginary part of (14) gives (4). He then showed, using the same partial fraction expansion as above, that

$$I_n = \frac{\pi i}{(2i)^n} \sum_{j=1}^n \left\{ A_j \sum_{\varepsilon \in \{-1, 1\}^n} \left( \prod_{r=1}^n \varepsilon_r \right) \operatorname{sgn} \left( \sum_{r=1}^n \varepsilon_r k_r \right) \exp \left( i \sum_{r=1}^n \varepsilon_r k_r (a_j - a_r) \right) \right\} \quad (15)$$

where  $a_1, a_2, \dots, a_n$  are distinct and

$$A_j := \frac{1}{\prod_{r \neq j} (a_j - a_r)}. \quad (16)$$

The formula (15) is quite elegant and also allows one to derive Corollary 2.6, which was independently found by Glasser in [7]. For instance, it suffices to appeal to the case  $m = 0$  of (11). However, as we will demonstrate in the remainder, the evaluation given in Theorem 2.4 has the advantage of making significant additional structure of the integrals  $I_n$  more apparent. Before doing so in Section 5 we next consider the case  $I_3$  in detail.

### 3 The case $n = 3$

We can completely dispose of the three-dimensional integral  $I_3$  by considering the three cases:  $a_1, a_2, a_3$  distinct;  $a_1$  distinct from  $a_2 = a_3$ ; and  $a_1 = a_2 = a_3$ .




Figure 2: Integrands in (1) with parameters  $\mathbf{a} = (-1, 0, 1)$  and  $\mathbf{k} = (k_1, 2, 1)$  where  $k_1 = 2, 4, 7$

### 3.1 The case $n = 3$ when $a_1, a_2, a_3$ are distinct

As demonstrated in this section, the evaluation of  $I_3$  will depend on which inequalities are satisfied by the parameters  $k_1, k_2, k_3$ . For  $n = 3$ , Theorem 2.4 yields:

$$I_3 \begin{pmatrix} a_1, a_2, a_3 \\ k_1, k_2, k_3 \end{pmatrix} = \frac{1}{8} \sum_{j=1}^3 \sum_{\varepsilon \in \{-1,1\}^3} \left[ \varepsilon_j \operatorname{sgn}(\varepsilon_1 k_1 + \varepsilon_2 k_2 + \varepsilon_3 k_3) \prod_{i \neq j} \frac{\sin(k_i(a_j - a_i))}{a_j - a_i} \right. \\ \left. - \varepsilon_1 \varepsilon_2 \varepsilon_3 \operatorname{sgn}(\varepsilon_1 k_1 + \varepsilon_2 k_2 + \varepsilon_3 k_3) \prod_{i \neq j} \frac{\cos(k_i(a_j - a_i))}{a_j - a_i} \right]. \quad (17)$$

**Remark 3.1** (Recovering Djokvič's Evaluation). Upon using the identity  $\sin(x)\sin(y) - \cos(x)\cos(y) = -\cos(x+y)$  to combine the two products, the right-hand side of equation (17) can be reexpressed in the symmetric form

$$-\frac{1}{8} \sum_{\varepsilon \in \{-1,1\}^3} \varepsilon_1 \varepsilon_2 \varepsilon_3 \operatorname{sgn}(\varepsilon_1 k_1 + \varepsilon_2 k_2 + \varepsilon_3 k_3) \sum_{j=1}^3 \frac{\cos(\sum_{i \neq j} \varepsilon_i k_i (a_j - a_i))}{\prod_{i \neq j} (a_j - a_i)}.$$

This is precisely Djokvič's evaluation (15). ◇

In fact, distinguishing between two cases, illustrated in Figure 3, the evaluation (17) of  $I_3$  can be made entirely explicit:

**Corollary 3.2** ( $a_1, a_2, a_3$  distinct). *Assume that  $k_1, k_2, k_3 > 0$ . Then*

1. *If  $\frac{1}{2} \sum k_i \leq k_\ell$ , as can happen for at most one index  $\ell$ , then:*

$$I_3 \begin{pmatrix} a_1, a_2, a_3 \\ k_1, k_2, k_3 \end{pmatrix} = \pi \prod_{i \neq \ell} \frac{\sin(k_i(a_\ell - a_i))}{a_\ell - a_i} \quad (18)$$

2. Otherwise, that is if  $\max k_i < \frac{1}{2} \sum k_i$ , then:

$$I_3 \begin{pmatrix} a_1, a_2, a_3 \\ k_1, k_2, k_3 \end{pmatrix} = \frac{\pi}{2} \sum_{j=1}^3 \left[ \prod_{i \neq j} \frac{\sin(k_i(a_j - a_i))}{a_j - a_i} + \prod_{i \neq j} \frac{\cos(k_i(a_j - a_i))}{a_j - a_i} \right] \quad (19)$$




Figure 3: The constraints on the parameters  $k_j$  in Corollary 3.2 — if  $k_1, k_2$  take values in the shaded regions then (18) holds with the indicated choice of  $\ell$

*Proof.* The first case is a special case of Corollary 2.6. Alternatively, assuming without loss that the inequality for  $k_\ell$  is strict, it follows directly from (17): because  $\text{sgn}(\varepsilon_1 k_1 + \varepsilon_2 k_2 + \varepsilon_3 k_3) = \varepsilon_\ell$  all but one sum over  $\varepsilon \in \{-1, 1\}^3$  cancel to zero.

In the second case,  $k_1 < k_2 + k_3$ ,  $k_2 < k_3 + k_1$ ,  $k_3 < k_1 + k_2$ . Therefore:

$$\begin{aligned} \frac{1}{8} \sum_{\varepsilon \in \{-1, 1\}^3} \varepsilon_j \text{sgn}(\varepsilon_1 k_1 + \varepsilon_2 k_2 + \varepsilon_3 k_3) &= \frac{1}{2} \quad \text{for all } j, \\ -\frac{1}{8} \sum_{\varepsilon \in \{-1, 1\}^3} \varepsilon_1 \varepsilon_2 \varepsilon_3 \text{sgn}(\varepsilon_1 k_1 + \varepsilon_2 k_2 + \varepsilon_3 k_3) &= \frac{1}{2}. \end{aligned}$$

The claim then follows from (17). □

**Remark 3.3** (Hidden Trigonometric Identities). Observe that because of the continuity of  $I_3$  as a function of  $k_1$ ,  $k_2$ , and  $k_3$ , we must have the non-obvious identity

$$\prod_{i \neq 1} \left[ \frac{\sin(k_i(a_1 - a_i))}{a_1 - a_i} \right] = \frac{1}{2} \sum_{j=1}^3 \left[ \prod_{i \neq j} \frac{\sin(k_i(a_j - a_i))}{a_j - a_i} + \prod_{i \neq j} \frac{\cos(k_i(a_j - a_i))}{a_j - a_i} \right] \quad (20)$$

when  $k_1 = k_2 + k_3$ . We record that *Mathematica 7* is able to verify (20); however, it struggles with the analogous identities arising for  $n \geq 4$ . ◇

### 3.2 The case $n = 3$ when $a_1 \neq a_2 = a_3$

As a limiting case of Corollary 3.2 we obtain:

**Corollary 3.4** ( $a_1 \neq a_2 = a_3$ ). Assume that  $k_1, k_2, k_3 > 0$  and  $a_1 \neq a_2$ . Set  $a := a_2 - a_1$ .

1. If  $k_1 \geq \frac{1}{2} \sum k_i$  then:

$$I_3 \left( \begin{matrix} a_1, a_2, a_2 \\ k_1, k_2, k_3 \end{matrix} \right) = \pi \frac{\sin(k_2 a)}{a} \frac{\sin(k_3 a)}{a} \quad (21)$$

2. If  $\max(k_2, k_3) \geq \frac{1}{2} \sum k_i$  then:

$$I_3 \left( \begin{matrix} a_1, a_2, a_2 \\ k_1, k_2, k_3 \end{matrix} \right) = \pi \min(k_2, k_3) \frac{\sin(k_1 a)}{a} \quad (22)$$

3. Otherwise, that is if  $\max k_i < \frac{1}{2} \sum k_i$ :

$$I_3 \left( \begin{matrix} a_1, a_2, a_2 \\ k_1, k_2, k_3 \end{matrix} \right) = \frac{\pi}{2} \frac{\cos((k_2 - k_3)a) - \cos(k_1 a)}{a^2} + \frac{\pi}{2} \frac{(k_2 + k_3 - k_1) \sin(k_1 a)}{a}. \quad (23)$$

*Proof.* The first two cases are immediate consequences of (18) upon taking the limit  $a_3 \rightarrow a_2$ .

Likewise, the third case follows from (19) with just a little bit of care. The contribution of the sine products from (19) is

$$\frac{\pi}{2} \frac{\sin(k_2 a) \sin(k_3 a)}{a^2} + \frac{\pi}{2} \frac{(k_2 + k_3) \sin(k_1 a)}{a}.$$

On the other hand, writing  $a_3 = a_2 + \varepsilon$  with the intent of letting  $\varepsilon \rightarrow 0$ , the cosine products contribute

$$\frac{\pi}{2} \left[ \frac{\cos(k_2 a) \cos(k_3 a)}{a^2} - \frac{\cos(k_1 a) \cos(k_3 \varepsilon)}{a \varepsilon} + \frac{\cos(k_1 (a + \varepsilon)) \cos(k_2 \varepsilon)}{(a + \varepsilon) \varepsilon} \right].$$

The claim therefore follows once we show

$$\frac{\cos(k_1 a) \cos(k_3 \varepsilon)}{a \varepsilon} - \frac{\cos(k_1(a + \varepsilon)) \cos(k_2 \varepsilon)}{(a + \varepsilon) \varepsilon} \rightarrow \frac{\cos(k_1 a)}{a^2} + \frac{k_1 \sin(k_1 a)}{a}.$$

This is easily verified by expanding the left-hand side in a Taylor series with respect to  $\varepsilon$ . In fact, all the steps in this proof can be done automatically using, for instance, *Mathematica* 7.  $\square$

Observe that, since  $I_n$  is invariant under changing the order of its arguments, Corollary 3.4 covers all cases where exactly two of the parameters  $a_j$  agree.

**Remark 3.5** (Alternative Approach). We remark that Corollary 3.4 can alternatively be proved in analogy with the proof given for Theorem 2.4 — that is by starting with a partial fraction decomposition and evaluating the occurring basic integrals. Besides integrals covered by equation (12) this includes formulae such as

$$\begin{aligned} & \int_{-\infty}^{\infty} \frac{\sin(k_2 x)}{x} \frac{\sin(k_3 x)}{x} \cos(k_1 x) dx \\ &= \frac{\pi}{8} \sum_{\varepsilon \in \{-1, 1\}^3} \varepsilon_2 \varepsilon_3 (\varepsilon_1 k_1 + \varepsilon_2 k_2 + \varepsilon_3 k_3) \operatorname{sgn}(\varepsilon_1 k_1 + \varepsilon_2 k_2 + \varepsilon_3 k_3). \end{aligned} \quad (24)$$

This evaluation follows from [3, Theorem 3(ii)]. In fact, (24) is an immediate consequence of equation (12) with  $n = 3$  and  $A = \emptyset$  after integrating with respect to one of the parameters  $k_i$  where  $i \in B$ . Clearly, this strategy evaluates a large class of integrals, similar to (24), over the real line with integrands products of sines and cosines as well as powers of the integration variable (see also [3]).  $\diamond$

### 3.3 The case $n = 3$ when $a_1 = a_2 = a_3$

In this case,

$$\begin{aligned} I_3 &= \int_{-\infty}^{\infty} \frac{\sin(k_1(x - a_1))}{x - a_1} \frac{\sin(k_2(x - a_1))}{x - a_1} \frac{\sin(k_3(x - a_1))}{x - a_1} dx \\ &= \int_{-\infty}^{\infty} \frac{\sin(k_1 x)}{x} \frac{\sin(k_2 x)}{x} \frac{\sin(k_3 x)}{x} dx. \end{aligned} \tag{25}$$

**Corollary 3.6** ( $a_1 = a_2 = a_3$ ). *Assume without loss that  $k_1 \geq k_2 \geq k_3 > 0$ . Then*

1. *If  $k_1 \geq k_2 + k_3$  then:*

$$I_3 \left( \begin{matrix} a_1, a_1, a_1 \\ k_1, k_2, k_3 \end{matrix} \right) = \pi k_2 k_3$$

2. *If  $k_1 \leq k_2 + k_3$  then:*

$$I_3 \left( \begin{matrix} a_1, a_1, a_1 \\ k_1, k_2, k_3 \end{matrix} \right) = \pi \left( k_2 k_3 - \frac{(k_2 + k_3 - k_1)^2}{4} \right)$$

*Proof.* The first part follows from Theorem 2 and the second from Corollary 1 in [3].

Alternatively, Corollary 3.6 may be derived from Corollary 3.4 on letting  $a$  tend to zero. Again, this can be automatically done in a computer algebra system such as *Mathematica 7* or *Maple 14*.  $\square$

## 4 Particularly special cases of sinc integrals

The same phenomenon as in Corollary 3.6 leads to one of the most striking examples in [3]. Consider the following example of a re-normalized  $I_n$  integral, in which we set:

$$J_n := \int_{-\infty}^{\infty} \operatorname{sinc} x \cdot \operatorname{sinc} \left( \frac{x}{3} \right) \cdots \operatorname{sinc} \left( \frac{x}{2n+1} \right) dx.$$

Then — as *Maple* and *Mathematica* are able to confirm — we have the following evaluations:

$$\begin{aligned} J_0 &= \int_{-\infty}^{\infty} \operatorname{sinc} x dx = \pi, \\ J_1 &= \int_{-\infty}^{\infty} \operatorname{sinc} x \cdot \operatorname{sinc} \left( \frac{x}{3} \right) dx = \pi, \\ &\vdots \\ J_6 &= \int_{-\infty}^{\infty} \operatorname{sinc} x \cdot \operatorname{sinc} \left( \frac{x}{3} \right) \cdots \operatorname{sinc} \left( \frac{x}{13} \right) dx = \pi. \end{aligned}$$

As explained in detail in [3] or [4, Chapter 2], the seemingly obvious pattern — a consequence of Corollary 2.6 — is then confounded by

$$\begin{aligned} J_7 &= \int_{-\infty}^{\infty} \operatorname{sinc} x \cdot \operatorname{sinc} \left( \frac{x}{3} \right) \cdots \operatorname{sinc} \left( \frac{x}{15} \right) dx \\ &= \frac{467807924713440738696537864469}{467807924720320453655260875000} \pi < \pi, \end{aligned}$$

where the fraction is approximately 0.9999999998529... which, depending on the precision of calculation, numerically might not even be distinguished from 1.

This is a consequence of the following general evaluation given in [3]:

**Theorem 4.1.** Denote  $K_m = k_0 + k_1 + \dots + k_m$ . If  $2k_j \geq k_n > 0$  for  $j = 0, 1, \dots, n-1$  and  $K_n > 2k_0 \geq K_{n-1}$  then

$$\int_{-\infty}^{\infty} \prod_{j=0}^n \frac{\sin(k_j x)}{x} dx = \pi k_1 k_2 \cdots k_n - \frac{\pi}{2^{n-1} n!} (K_n - 2k_0)^n. \quad (26)$$

Note that Theorem 4.1 is a “first-bite” extension of Corollary 2.6: assuming only that  $k_j > 0$  for  $j = 0, 1, \dots, n$  then if  $2k_0 > K_n$  the integral evaluates to  $\pi k_1 k_2 \cdots k_n$ .

Theorem 4.1 makes clear that the pattern that  $J_n = \pi$  for  $n = 0, 1, \dots, 6$  breaks for  $J_7$  because

$$\frac{1}{3} + \frac{1}{5} + \dots + \frac{1}{15} > 1$$

whereas all earlier partial sums are less than 1.

Yet, we do have the surprising equality [2] of the integrals  $J_n$  and corresponding Riemann sums:

$$\int_{-\infty}^{\infty} \prod_{j=0}^n \operatorname{sinc}\left(\frac{x}{2j+1}\right) dx = \sum_{m=-\infty}^{\infty} \prod_{j=0}^n \operatorname{sinc}\left(\frac{m}{2j+1}\right) \quad (27)$$

for  $n = 1, 2, \dots, 7, 8, \dots, 40248$ . For  $n > 40248$  this equality fails as well; the sum being strictly bigger than the integral. As in the case of (26) there is nothing special about the choice of parameters  $k_j = \frac{1}{2j+1}$  in the sinc functions [2]:

**Theorem 4.2.** Suppose that  $k_1, k_2, \dots, k_n > 0$ . If  $k_1 + k_2 + \dots + k_n < 2\pi$  then

$$\int_{-\infty}^{\infty} \prod_{j=1}^n \operatorname{sinc}(k_j x) dx = \sum_{m=-\infty}^{\infty} \prod_{j=1}^n \operatorname{sinc}(k_j m). \quad (28)$$

As a consequence, we see that (27) holds for  $n$  provided that

$$\sum_{j=0}^n \frac{1}{2j+1} < 2\pi$$

which is true precisely for the range of  $n$  specified above.

**Remark 4.3.** With this insight, it is not hard to contrive more persistent examples. An entertaining example given in [2] is taking the reciprocals of primes: using the Prime Number Theorem one estimates that the sinc integrals equal the sinc sums until the number of products is about  $10^{176}$ . That of course makes it rather unlikely to find by mere testing an example where the two are unequal. Even worse for the naive tester is the fact that the discrepancy between integral and sum is always less than  $10^{-10^{86}}$  (and even smaller if the Riemann hypothesis is true).  $\diamond$

A related integral which because of its varied applications has appeared repeatedly in the literature, see e.g. [9] and the references therein, is

$$\frac{2}{\pi} \int_0^\infty \left( \frac{\sin x}{x} \right)^n \cos(bx) dx \quad (29)$$

which, for  $0 \leq b < n$ , has the closed form

$$\frac{1}{2^{n-1}(n-1)!} \sum_{0 \leq k < (n+b)/2} (-1)^k \binom{n}{k} (n+b-2k)^{n-1}.$$

To give an idea of the range of applications, we only note that the authors of [9] considered the integral (29) because it is proportional to “the intermodulation distortion generated by taking the  $n$ th power of a narrow-band, high-frequency white noise”; on the other hand, the recent [1] uses (29) with  $b = 0$  to obtain an improved lower bound for the Erdős-Moser problem.

If  $b \geq n$  then the integral (29) vanishes. The case  $b = 0$  in (29) is the interesting special case of  $I_n$  with  $k_1 = \dots = k_n = 1$  and  $a_1, \dots, a_n = 0$ . Its evaluation appears as an exercise in [12, p. 123]; in [3] it is demonstrated how it may be derived using the present methods.

## 5 The case $n \geq 4$

Returning to Theorem 2.4 we now show that in certain general cases the evaluation of the integral  $I_n$  may in essence be reduced to the evaluation of the integral  $I_m$  for some  $m < n$ . In particular, we will see that Corollary 2.6 is the most basic such case — corresponding to  $m = 1$ .

In order to exhibit this general structure of the integrals  $I_n$ , we introduce the notation

$$I_{n,j} := \sum_{A,B} \alpha_{j,A,B} C_{j,A,B} \quad (30)$$

so that, by (13),  $I_n = \sum_{j=1}^n I_{n,j}$ .

**Theorem 5.1** (Substructure). *Assume that  $k_1 \geq k_2 \geq \dots \geq k_n > 0$ , and that  $a_1, a_2, \dots, a_n$  are distinct. Suppose that there is some  $m$  such that for all  $\varepsilon \in \{-1, 1\}^n$  we have*

$$\operatorname{sgn}(\varepsilon_1 k_1 + \dots + \varepsilon_m k_m + \dots + \varepsilon_n k_n) = \operatorname{sgn}(\varepsilon_1 k_1 + \dots + \varepsilon_m k_m). \quad (31)$$

Then

$$I_n = \sum_{j=1}^m I_{m,j} \prod_{i>m} \frac{\sin(k_i(a_j - a_i))}{a_j - a_i}. \quad (32)$$

*Proof.* Note that in light of (12) and (31) we have  $\alpha_{j,A,B} = 0$  unless  $\{m+1, \dots, n\} \subset B$ . To see this assume that there is some index  $k > m$  such that  $k \in A$  or  $k = j$ . Then the term in (12) contributed by  $\varepsilon \in \{-1, 1\}^n$  has opposite sign as the term contributed by  $\varepsilon'$ , where  $\varepsilon'$  is obtained from  $\varepsilon$  by flipping the sign of  $\varepsilon_k$ . The claim now follows from Theorem 2.4.  $\square$

**Remark 5.2.** The condition (31) may equivalently be stated as

$$\min |\varepsilon_1 k_1 + \dots + \varepsilon_m k_m| > k_{m+1} + \dots + k_n \quad (33)$$

where the minimum is taken over  $\varepsilon \in \{-1, 1\}^m$ . We idly remark that, for large  $m$ , computing the minimum is a hard problem. In fact, in the special case of integral  $k_j$  just deciding whether the minimum is zero (which is equivalent to the *partition problem* of deciding whether the parameters  $k_j$  can be partitioned into two sets with the same sum) is well-known to be NP-complete [6, Section 3.1.5].  $\diamond$

Observe that the case  $m = 1$  of Theorem 5.1 together with the basic evaluation (4) immediately implies Corollary 2.6. This is because the condition (31) holds for  $m = 1$  precisely if  $k_1 > k_2 + \dots + k_n$ .

If (31) holds for  $m = 2$  then it actually holds for  $m = 1$  provided that the assumed inequality  $k_1 \geq k_2$  is strict. Therefore the next interesting case is  $m = 3$ . The final evaluation makes this case explicit. It follows from Corollary 3.2.

**Corollary 5.3** (A second n-dimensional case). *Let  $n \geq 3$ . Assume that  $k_1 \geq k_2 \geq \dots \geq k_n > 0$ , and that  $a_1, a_2, \dots, a_n$  are distinct. If*

$$k_1 \leq k_2 + \dots + k_n \text{ and } k_2 + k_3 - k_1 \geq k_4 + \dots + k_n$$

then:

$$I_n = \frac{\pi}{2} \sum_{j=1}^3 \prod_{i \geq 4} \frac{\sin(k_i(a_j - a_i))}{a_j - a_i} \left[ \prod_{i < 3, i \neq j} \frac{\sin(k_i(a_j - a_i))}{a_j - a_i} + \prod_{i \leq 3, i \neq j} \frac{\cos(k_i(a_j - a_i))}{a_j - a_i} \right].$$

The cases  $m \geq 4$  quickly become much more involved. In particular, the condition (31) becomes a set of inequalities.

To close, we illustrate with the first case not covered by Corollaries 2.6 and 5.3:

**Example 5.4.** As usual, assume that  $k_1 \geq k_2 \geq k_3 \geq k_4 > 0$ , and that  $a_1, a_2, a_3, a_4$  are distinct. If  $k_1 < k_2 + k_3 + k_4$  (hence Corollary 2.6 does not apply) and  $k_1 + k_4 > k_2 + k_3$  (hence Corollary 5.3 does not apply either) then

$$\begin{aligned}
 I_4 &= \frac{\pi}{4} \sum_{j=1}^4 \sum_{A,B} C_{j,A,B} + \frac{\pi}{2} \prod_{i \neq 1} \frac{\sin(k_i(a_1 - a_i))}{a_1 - a_i} \\
 &\quad - \frac{\pi}{2} \sum_{j=2}^4 \frac{\sin(k_1(a_j - a_1))}{a_j - a_1} \prod_{i \neq 1, j} \frac{\cos(k_i(a_j - a_i))}{a_j - a_i}
 \end{aligned} \tag{34}$$

where the summation in the first sum is as in Theorem 2.4. Note that the terms  $I_{4,j}$  of (30) are implicit in (34) and may be used to make the case  $m = 4$  of Theorem 5.1 explicit as has been done in Corollary 5.3 for  $m = 3$ .  $\diamond$

## 6 Conclusions

We present these results for several reasons. First, the forensic nature of the mathematics was entertaining. Second, it made us reflect on how computer packages and databases have changed mathematics over the past forty to fifty years. Finally some of the evaluations merit being better known as they are excellent tests of computer algebra or numerical integration.

**Acknowledgments.** We want to thank Larry Glasser for pointing us to this problem after hearing a lecture by the first author on [3] and for providing historic context. We are also thankful for his and Tewodros Amdeberhan's comments on an earlier draft of this manuscript.

# References

- [1] Iskander Aliev. Siegel's lemma and Sum-Distinct sets. *Discrete Comput. Geom.*, 39(1-3):59–66, 2008.
- [2] Robert Baillie, David Borwein, and Jonathan Borwein. Surprising sinc sums and integrals. *This MONTHLY*, 115(10):888–901, 2008.
- [3] David Borwein and Jonathan Borwein. Some remarkable properties of sinc and related integrals. *The Ramanujan Journal*, 5:73–90, 2001.
- [4] J. M. Borwein, D. H. Bailey, and R. Girgensohn. *Experimentation in Mathematics: Computational Paths to Discovery*. A. K. Peters, 1st edition, 2004.
- [5] Timothy Feeman. *A Beginner's Guide to The Mathematics of Medical Imaging*. Springer Undergraduate Texts in Mathematics and Technology, 2010.
- [6] Michael R. Garey and David S. Johnson. *Computers and Intractability: A Guide to the Theory of NP-Completeness*. W. H. Freeman and Company, 1979.
- [7] M. L. Glasser. Private communication. June 2011.
- [8] M. L. Glasser, William A. Newcomb, and A. A. Jagers. A multiple sine integral. *This MONTHLY*, 94(1):83–86, 1987.
- [9] R. G. Medhurst and J. H. Roberts. Evaluation of the integral  $I_n(b) = \frac{2}{\pi} \int_0^\infty \left(\frac{\sin x}{x}\right)^n \cos(bx) dx$ . *Mathematics of Computation*, 19(89):113–117, 1965.
- [10] Kent E. Morrison. Cosine products, fourier transforms, and random sums. *This MONTHLY*, 102(8):716–724, 1995.
- [11] Byron Schmuland. Random harmonic series. *This MONTHLY*, 110(5):407–416, 2003.
- [12] E. T. Whittaker and G. N. Watson. *A Course of Modern Analysis*. Cambridge University Press, 4th edition, 1927.

• Preprints of references [2] and [3] are available at <http://docserver.carma.newcastle.edu.au/>.